

Yield trial and sensory evaluation of sweetpotato cultivars in Highland Papua and West Papua Indonesia

by Saraswati Prabawardani

Submission date: 08-May-2023 06:01PM (UTC+0700)

Submission ID: 2087446496

File name: Yield_Trial_and_Sensory_Evaluation_of_Sweetpotato.pdf (1.02M)

Word count: 5302

Character count: 27930

1 Yield trial and sensory evaluation of sweetpotato cultivars in Highland Papua and West Papua Indonesia

P. Saraswati¹, A. Soplanit², A. T. Syahputra³, L. Kossay³, N. Muid³, E. Ginting⁴ and G. Lyons^{5*}
¹Faculty of Agriculture Papua State University Manokwari, West Papua; ²Assessment Institute for Agricultural Technology, Papua; ³International Potato Center, Bogor; ⁴Research Institute for Legume and Tuber Crops, Malang; ⁵University of Adelaide, South Australia.

16 Received 25 May 2013; received in revised form 15 August 2013; accepted 22 September 2013

Abstract

With the aim of improving income and nutrition of indigenous highland people in Papua and West Papua provinces of Indonesia, trials involving introduced and local sweetpotato (*Ipomoea batatas* L.) cultivars were conducted in the Baliem Valley and Arfak Mountains (Minyambouw), Indonesia. Tuber yield and other agronomic traits, chemical content and sensory traits for 17 introduced and three local sweetpotato cultivars were evaluated. Total tuber yield varied from 7.4 to 20.8 Mg ha⁻¹ in the Baliem Valley and from 1.9 to 14.4 Mg ha⁻¹ in Minyambouw. The introduced cultivars produced higher yields than the local cultivars in both regions. Cultivar Papua Pattipi produced the highest yield in the Baliem Valley, while Sawentar produced the highest yield in Minyambouw. Cultivars varied in dry matter, protein, and beta-carotene. Worembai was best for tuber flesh colour, taste, texture and sweetness in the Baliem Valley, while Helaleke and Papua Salosa scored highest for all sensory traits in Minyambouw. Introduced sweetpotato cultivars adapted well to both areas. In order to reduce malnutrition and increase income, it is important to grow several sweetpotato cultivars in each food garden, including high yielding cultivars and those with specific nutritional traits, such as high protein, provitamin A and anthocyanins.

Keywords: Sweetpotato, *Ipomoea batatas* (L.), Adaptation, Tuber yield, Sensory evaluation.

Introduction

26 Sweetpotato is an important staple food crop in the eastern part of Indonesia, particularly on the island of Papua. It is the main source of carbohydrate intake for the highlanders of Papua, where it accounts for around 95 % of the diet (Ruinard, 1969; Schneider et al., 1993). Sweetpotato and pigs are an integral part of the social and cultural system of indigenous communities.

In Minyambouw, sweetpotato is usually grown along with other root crops and vegetables without any agronomic input in the mixed cropping system. On the Baliem Valley floor it is cultivated, in both

23 monoculture and mixed cropping systems, in mounds on large, raised beds with a developed drainage system (Peters, 2001), and is grown all year round in both regions.

3 Intensive research efforts to enhance production and consumption have been undertaken in recent decades with the aim to explore the potential of sweetpotato as a nutritious food for humans and animals (Woolfe, 1992; Yamakawa and Yoshimoto, 2002). Sweetpotato tubers are an excellent source of energy, vitamins A (in orange tuber flesh cultivars), B and C, potassium, phosphorus, calcium and magnesium (Hill, 1984; Woolfe, 1992). Sweetpotato can produce more edible energy per

*Author for correspondences: Phone (08) 83136533; Email<graham.lyons@adelaide.edu.au>.

hectare per day than other food crops in the tropics (Woolfe, 1992). Orange-fleshed varieties provide beta-carotene, which acts as a precursor of vitamin A (Ofori et al., 2009), while purple-fleshed sweetpotatoes are rich in anthocyanin and have high antioxidant activity (Lila, 2004; Kano et al., 2005). Young sweetpotato leaves are rich in polyphenols, protein, vitamins and minerals and are used as a green vegetable (Islam et al., 2002; Johnson and Pace, 2010). Protein content of sweetpotato leaf ranges from 25.5-29.8% (An et al., 2003), which is higher than the protein of sweetpotato tubers, which ranges between 1.4 and 9.4% (Oboh et al., 1989).

Two research projects under the auspices of the Australian Centre for International Agricultural Research (ACIAR) have aimed to improve the livelihood of indigenous highland people of Papua and West Papua by increasing income and nutritional status through improving sweetpotato quality and production and pig management. The first project (2000-2008) was implemented in the Baliem Valley and the second was extended to a highland area (Minyambouw) of West Papua Province. A baseline survey conducted prior to the research reported that the average family in the Arfak mountain region has just one child under 15 years, and there are few people over 45 years old. These conditions are similar to those in the Baliem Valley, where a similar survey was conducted in the first phase of the ACIAR project (S. Mahalaya, pers. comm., 2011). The most widely grown and preferred sweetpotato in these areas are white or cream-fleshed cultivars, without beta-carotene and anthocyanin. Inadequate nutrients in the diet, in particular protein and those minerals and vitamins associated with protein (iron, zinc, selenium, vitamin A), is likely to be a major factor in the poor health indices observed in these areas.

Even though sweetpotato is the major staple food, most of the local sweetpotato clones are low yielding, with average yield of 5-10 Mg ha⁻¹ in the Baliem Valley (Peters, 2001) and 5 Mg ha⁻¹ in

Minyambouw (Syahputra et al., pers. comm. 2009). Introduction and selection of advanced sweetpotato cultivars for human food and pig feed may provide higher yields, protein and starch content and improve nutrition. Field trials are needed to assess the adaptability of improved cultivars, including Pattipi, Salosa and Sawentar, to local conditions. The adaptation trials of sweetpotato cultivars for human food and pig feed in the current study were conducted at highland sites in Papua and West Papua in 2010, with the aim to compare tuber yield, dry matter content, nutrient content and sensory traits (including taste, texture and sweetness). Improved sweetpotato production, especially when combined with suitable agroforestry programs, has a strong potential to improve food security in South-East Asia (Kumar, 2006).

Materials and Methods

Study sites

The study was conducted in collaboration with local farmers in two highland sites, the Baliem Valley in Papua Province and Minyambouw in West Papua Province. All sweetpotato cultivars were grown in farmers' gardens. Climate and soil conditions of both sites are presented in Table 1.

The amount of rainfall in the Baliem Valley between the first and third months after planting was an average of 271 mm/month, but declined from the fourth month until harvest time to an average of 50 mm of rainfall/month. The rainfall in Minyambouw was around a mean 260 mm/month (range 143-401 mm/month) during the trial period.

Sweetpotato cultivars

Twenty sweetpotato cultivars with important characteristics, such as high yield, short maturity, superior nutrition and high dry matter were collected from research institutions and both local highland sites (Table 2).

Table 1. Climate and soil conditions in the Baliem Valley and Minyambouw.

	Baliem Valley	Minyambouw
Latitude	138°30'-139°40' E; 3°45'-4°20'S	133°52'-133°54' E and 01°06'-01°08'
Altitude	1500-2000 m	1650-1750 m
Annual rainfall	1907 mm	2283 mm
Relative humidity	80%	75-85 %
Minimum temperature	15 °C	-
Maximum temperature	26 °C	-
Average temperature	19.3 °C	-
Solar radiation	1.38 kJ/cm ¹²	-
Daily sunlight duration	3.98 hours/day ³	-
Soil pH (H ₂ O)	5.0-6.0	5.4- 6.3
Soil type	sandy clay loam to silty clay with block structure	sandy clay loam to clay loam
Soil P (Bray method)	7 µg/g (low)	6 µg/g (low)
Soil N total (Kjeldahl method)	0.12-0.24 % (low to medium)	0.48 % (medium)
Soil C (Calorimetric method)	2-2.75 % (low to medium)	2.39 % (medium)
Soil K (Colwell method)	Low ²	Low ²

Source: ¹Peters, 2001; ²Schroo, 1963 in Soenarto and Rumawas, 1997; ³Unipa, 2011, unpublished data.



36

Figure 1. Map of Papua showing the areas where the sweetpotato trials were conducted

5

Experimental design

The field trials were arranged in a randomised complete block design (RCBD) with three replications at each site. The size of each plot was 3 m x 3 m. Spacing between mounds was 1 m. There

were nine mounds, each planted with two sweetpotato tip cuttings (25-30 cm length). Each mound was 40 cm high and 100 cm in diameter. Weeding was conducted at two, three and five months after planting. Sweetpotato cuttings were planted in March 2010 and harvested seven months

Table 2. Origin and tuber morphological characteristics of 20 sweetpotato cultivars.

No	Cultivar/clone	Origin	Skin colour	Flesh colour
1	Cangkuang	Balitkabi ¹	Dark red	Light yellow
2	Sawentar	Balitkabi	Red	Yellow
3	Sukuh	Balitkabi	Yellow	Light orange
4	Sari	Balitkabi	Red	Dark yellow
5	Papua Salosa	Balitkabi	Yellow	Yellow with orange spots
6	Papua Pattipi	Balitkabi	Light yellow	Light yellow
7	BB-20413-13	CIP ²	Light yellow	White
8	BB-00105-10	CIP	Orange	Dark orange
9	Ayamurasaki	CIP	Dark red	Purple
10	UnguFiriange ³	CIPUnipa	Dark redDark red	Dark purpleLight yellow
11	Airani-3	PPUS ⁴	Yellow/Cream	Light yellow
12	Miencon	PPUS	Light red (slightly pink)	Light orange
13	Worembai	PPUS	Red	Pale yellow
14	Dosak-1	PPUS	Dark red purplish	Slightly dark purple
15	Dosak-2	PPUS	Yellow slightly cream	Light yellow, purple spots
16	Numfor- 5	PPUS	Light yellow, red spot	Pale yellow
17	Nabire	PPUS	Red	White
18	Musan	B. Valley ⁵	Light red	Pale yellow
19	Helaleke	B. Valley	Red	Light yellow
20	Bramwamrum	Minyambouw ⁶	Dark red	Light yellow

¹ Balitkabi (Root Crops and Legumes Research Centre, Malang), ² CIP (International Potato Center, Bogor), ³ Instead of cultivar Firiange which was tested in Minyambouw, cultivar Ungu was used in the Baliem Valley, as Firiange did not grow there. ⁴ PPUS (Root Crop and Sago Research Centre, Manokwari), ⁵ Local Baliem Valley cultivars, ⁶ Local Minyambouw cultivar.

after planting in the Baliem Valley, and nine months after planting in Minyambouw, following consultation with local farmers. Plant leaf samples were collected four months after planting from each site and selected minerals analysed using inductively coupled plasma optical emission spectrometry (ICP-OES) to determine nutrient content.

Total tuber yield and dry matter content were determined after harvesting the tubers. Nutrient contents of samples were determined using the AOAC (1990) method. Protein was analysed by the micro-Kjeldahl procedure. Protein concentration was calculated using the following formula: $N (\%) = (\text{ml HCl} - \text{ml blank}) \times N \times 14.007 \times 100$. Protein (% dry weight) = % N x conversion factor (6.25). Fat content was analysed using the Soxhlet method. Carbohydrate comprised the remainder of the sample. Beta-carotene was analysed using the

HPLC method described by Rodriguez-Amaya and Kimura (2004).

Evaluation of sensory traits was conducted by semi-trained consumer panels comprising farmers, teachers and housewives at both sites. Tuber samples of the 20 sweetpotato cultivars were placed in labelled plastic bags, and steamed for 30 minutes. The cooked samples were served to panellists randomly and evaluated for colour acceptability, taste, texture, fibrosity, sweetness level, performance and general acceptability. The responses of the panellists were recorded on a five-point hedonic scale. The panellists rinsed their mouth with plain water after tasting each sample. The agronomic data collected from the sites were analysed by standard ANOVA using GenStat version 14 at a 95% confidence interval, while sensory characteristics were analysed descriptively.

Results and Discussion

14

Tuber yield and dry matter content

Sweetpotato cultivars varied significantly in tuber yield and dry matter content ($P < 0.05$). Average tuber yield was higher in the Baliem Valley (average of 12.1 Mg ha^{-1}) than in Minyambouw (average of 9.6 Mg ha^{-1}) (Fig. 2). Yield of sweetpotato cultivars in the Baliem Valley ranged from 7.4 Mg ha^{-1} to 20.8 Mg ha^{-1} , with the highest yielding (20.8 Mg ha^{-1}) being the introduced cultivar Papua Pattipi. Other relatively high yielding cultivars (with more than 12 Mg ha^{-1}) were Musan (a local Baliem Valley cultivar), Papua Salosa, Sawentar, Dosak-2, BB-20413.13, Ungu, Cangkuang, Helaleke (local) and Numfor-5.

Tuber yield varied according with the location. In Minyambouw, the average tuber yield was highest in cultivar Sawentar, which produced 14.4 Mg ha^{-1} . Other relatively high yielding cultivars ($10.1\text{--}13.9 \text{ Mg ha}^{-1}$) were Airani-3, Numfor-5, Miencon, BB-20413-13, Worembai, Helaleke, Papua Pattipi, Dosak-1 and Papua Salosa. The local Minyambouw cultivar (Bramwamrum) produced 7.5 Mg ha^{-1} . Firiange and Ayamurasaki produced the lowest yields, 4 Mg ha^{-1} and 1.9 Mg ha^{-1} , respectively.

Bramwamrum produced the highest dry matter content in the Baliem Valley (36.5%) and in Minyambouw (39.6%), while the lowest dry matter was recorded from cultivar BB-00105.1 in both sites; 20.4% in the Baliem Valley and 18.3% in Minyambouw. Figure 2 shows that most cultivars yielded lower dry matter content in Minyambouw than in the Baliem Valley, except for the local Bramwamrum cultivar, which contained greater dry matter when it was planted in Minyambouw than in the Baliem Valley.

Tuber yield varied according to both cultivar and location. Several introduced cultivars produced higher yields than the local cultivars in both sites. The Baliem Valley and Minyambouw differ in




Figure 2. Yield and dry matter content of tubers collected from (a) the Baliem Valley and (b) Minyambouw sites. Note that the y axis scales vary.

Baliem Valley: yield CV%: 18.3, LSD (0.05): 3.7; dry matter CV%: 9.3, LSD: 4.9

Minyambouw: yield CV%: 30.3, LSD: 4.8; dry matter CV%: 5.4, LSD: 2.5.

climate, topography and soil type. Sweetpotato leaves from the Baliem Valley had higher boron, potassium and sulphur concentration for the same cultivar than those from Minyambouw (data not shown). This may have contributed to the higher tuber yield and higher dry matter content of the tested sweetpotato cultivars in the Baliem Valley.

The yield of tubers grown in the Baliem Valley in the present study was 28% lower than that in the previous trial of the ACTAR phase I project (Jusuf et al., 2007; Soplanit et al., 2005). This can be attributed to the dry conditions due to La Niña that

occurred in the Baliem Valley from the time of tuber bulking until harvest. However, most of the cultivars were well adapted to the local environment, as shown by production of more than 10 Mg ha⁻¹, which is higher than the mean national sweetpotato yield of 10 Mg ha⁻¹ (International Potato Center, 1998) and that recorded for the Baliem Valley of 5-10 Mg ha⁻¹ (Peters, 2001).

Sweetpotato normally yields 5 Mg ha⁻¹ in Minyambouw (Syahputra et al., pers. comm., 2009). In this area, it is traditionally cultivated without any soil tillage. Application of appropriate methods of cultivation, such as mounding and optimal plant spacing increased yield in this trial. Yield of the local cultivar Bramwamrum increased from the usual level of 5 Mg ha⁻¹ to 7 Mg ha⁻¹. Mounding allows more oxygen circulation and improved drainage for tuber development, and facilitates harvesting (Sillitoe, 1998).

Local varieties normally take nine to ten months to mature in high altitude areas. However, the introduced cultivars in this study matured earlier, taking just six months from planting to harvest in the Baliem Valley. These early maturing cultivars offer an opportunity for faster recovery of sweetpotato yield after drought, which periodically occurs in the Baliem Valley. Some introduced cultivars in the Minyambouw trial, namely Papua Pattipi, Papua Salosa, Cangkuang and Sawentar,

were also visually observed to be ready for harvest at seven months after planting. This could be important for food security.

Nutrient Analysis

Tuber nutrient content varied significantly among the 20 sweetpotato cultivars (Table 3). Protein content was highest in cultivars Ungu, Ayamurasaki, Papua Salosa and Numfor-5, which contained 4.6%, 4.5%, 4.5% and 4.4%, respectively. The local Minyambouw cultivar (Bramwamrum), which had the highest dry matter content, also had the greatest starch content of 37.2%, followed by Papua Salosa (35.4%), and Papua Pattipi (35.3%). The orange tuber flesh cultivar BB-00105.10, which had the lowest dry matter content, also had the lowest starch content (16.7%). The fat content of all cultivars was low, varying from 0.1 to 0.8%; Sawentar had the highest fat content (0.8%). Cultivar Nabire had the highest ash content (2.5%), followed by Helaleke (1.4%), Dosak-2 (1.3%) and Dosak 1 (1.3%).

Nutrients of the 20 sweetpotato cultivars were analyzed using tubers harvested from the Baliem Valley trial (Table 3). The Minyambouw data were not included due to a faulty analytical instrument. Differences in yield and nutrient concentrations are attributed to genotype-environment interaction. Papua Pattipi had higher iron, zinc, manganese, copper and calcium, but lower potassium and

Table 3. Mean nutrient contents of sweetpotato cultivars grown in the Baliem Valley. Data for the highest five and lowest one for protein are shown of the 20 sweetpotato cultivars grown.

Sample	Protein (%)	Carbohydrate (%)	Fat (%)	Ash (%)
Ungu/Firiangge	4.55	26.47	1.07	1.07
Ayamurasaki	4.54	30.37	0.97	0.97
Papua Salosa	4.52	35.37	0.67	0.67
Numfor-5	4.35	27.89	0.57	0.57
Worembai	3.91	31.85	1.08	1.08
Sari	1.20	20.59	0.32	0.32
Mean	3.09	28.68	0.93	0.93
CV(%)	3.4	0.90	2.50	2.50
LSD (P=0.05)	0.22**	0.52**	0.05**	0.05**

Mean, CV and LSD for all 20 cultivars grown.

phosphorus in tubers than Bramwamrum grown at the same site (data not shown). The lower potassium and phosphorus in Papua Pattipi are attributed to the dilution of the minerals by the formation of heavier tubers.

Improved nutritional content of sweetpotato tubers can reduce malnutrition and improve food security in the developing world. It was interesting that two of the purple-fleshed cultivars contained the highest protein of all the cultivars: Ungu (4.6%) and Ayamurasaki (4.5%), followed by two yellow-fleshed cultivars, Papua Salosa (4.5%) and Numfor-5 (4.4%). The lowest protein was in the cultivars Bramwamrum (1.6%) and Sari (1.2%). The protein content of sweetpotato in other studies included a 0.5-5% range in Papua New Guinea (Bradbury et al., 1984).

Beta-carotene was analysed in selected sweetpotato cultivars that represented the yellow and orange tuber flesh colour. The cultivar with deep orange tuber flesh colour (BB-00105-10) had the highest beta-carotene level (226 $\mu\text{g g}^{-1}$ dry weight). The pale orange or yellow-fleshed cultivars contained much lower amounts of beta-carotene (average of 20 $\mu\text{g g}^{-1}$) (Table 4). Beauregard, an orange-fleshed cultivar which is common in developed countries was reported to have 215 $\mu\text{g g}^{-1}$ dry weight of beta-carotene (Ofori et al., 2009). The intensity of the sweetpotato's orange flesh colour is directly related to its beta-carotene content, and some dark orange cultivars contain similar levels to carrots (Woolfe, 1992; Takahata et al., 1993). Cultivar BB-00105.10 could therefore play an important role in reducing vitamin A deficiency.

Sensory analysis

The local panellists' preferences differed between the two regions. In the Baliem Valley, Worembai scored the highest mean value (4.5) for all sensory attributes (Table 5). In this area, the purple-fleshed cultivars Ayamurasaki, Dosak-1 and Ungu were among the five cultivars scored by the local people

Table 4. Beta-carotene levels of orange- and yellow-fleshed tuber sweetpotato cultivars.

Cultivar	Flesh colour	Beta-carotene ($\mu\text{g/g DW}$)
BB-00105-190	Dark orange	226
Sukuh	Pale orange	40
Papua Salosa	Yellow/orange spots	35
Miencon	Pale orange/cream	12
Sawentar	Yellow	10
Sari	Yellow	5
Nabire	White	0

as having the best taste. However, BB-00105.10, which is the only cultivar with dark orange flesh colour, and which contained the highest beta-carotene level, was less preferred by the panellists, scoring 3.7 due to poor texture and high water content. Consumers in the Baliem Valley also considered Bramwamrum, Ayamurasaki, Ungu, Dosak-1, Papua Pattipi, Sawentar, Papua Salosa, Cangkuang and Helaleke to have desirable sensory traits, with scores between 4.2 and 3.8 (Table 5).

On the other hand, Helaleke and Papua Salosa were the most preferred cultivars by Minyambouw's panellists, both scoring 4.9, followed by Sukuh, Sawentar, Dosak-2, Sari and Cangkuang, scoring between 4.7 and 4.3 (Table 6). A number of cultivars (Firiange, Miencon, Dosak-1, BB-20413-13 and BB-00105.10) received low scores (between 2.6 and 2.9) in Minyambouw.

Most of the tested cultivars were found to have acceptable tuber flesh colour, except for Musan in both sites. Musan was the least preferred cultivar for human consumption with low scores in all sensory characters, due to poor texture and high fibre. This cultivar produced a high yield in the Baliem Valley but the overall sensory characters showed that it was not regarded as fit for staple consumption by local people. It is usually fed to their domestic animals, particularly pigs.

Panellists from both areas considered that high dry matter content (also expressed as starchy) and good

Table 5. Sensory scores for steam cooked tubers of sweetpotato cultivars in the Baliem Valley, using a hedonic scale¹. Data included from the six highest-scoring cultivars and the lowest cultivar for acceptability and taste of the 20 tested.

Cultivar	Colour acceptability	Taste	Texture	Fibre	Sweetness level	Appearance	General acceptability	Mean
Bramwamrum	4.1	5.0	4.0	3.8	3.6	3.9	5.0	4.2
Worembai	4.7	4.7	4.7	4.0	4.1	4.0	5.0	4.5
Ayamurasaki	4.1	4.5	4.0	3.9	3.3	4.0	5.0	4.1
Dosak-1	4.2	4.3	3.6	3.7	3.8	4.1	5.0	4.1
Ungu	4.3	4.2	3.9	4.0	3.2	4.2	5.0	4.1
Papua Salosa	4.3	4.1	3.3	4.0	3.7	3.8	5.0	4.0
Musan	2.8	2.6	2.3	2.2	3.2	2.9	2.0	2.6

¹Five-point hedonic scale from 1 = dislike to 5 = extremely like.

Table 6. Sensory scores for steam-cooked tubers of sweetpotato cultivars in Minyambouw using a hedonic scale¹. Data included from the six highest-scoring cultivars and the lowest cultivar for acceptability and taste of the 20 tested.

Cultivar	Colour acceptability	Taste	Texture	Fibre	Sweetness level	Appearance	General acceptability	Mean
Helaleke	4	5	5	5	5	5	5	4.9
Papua Salosa	4	5	5	5	5	5	5	4.9
Sukuh	4	5	4	5	5	5	5	4.7
Sawentar	4	5	5	4	5	5	5	4.7
Dosak-2	3	5	5	4	5	4	5	4.4
Sari	3	5	3	5	5	5	5	4.4
Dosak-1	4	1	1	5	1	4	3	2.7

¹Five-point hedonic scale from 1 = dislike to 5 = extremely like.

taste were the most important criteria. Dry matter content depends on cultivar, location, climate, daylength, pathogens and cultivation practices (Woolfe, 1992). For example, high humidity as a result of heavy rainfall during the trial in Minyambouw reduced tuber dryness. According to Rahman et al. (2003), genotypes with high starch content have lower tuber water content when harvested in the dry season. A sensory test of sweetpotato in Kenya also showed an adult preference for sweetpotato cultivars with dry matter greater than 27% (Hagenimana et al. 2001). The average dry matter content in this trial was 31.9% in the Baliem Valley and 27.5% in Minyambouw, which is in broad agreement with the mean of 30% reported by Ekayanake and Collins (2004).

Some cultivars, including the purple-fleshed Ayamurasaki, Dosak-1 and Ungu, were favoured

by panellists in both regions. These are known to contain high antioxidant activity due mainly to their high level of anthocyanins (Teow et al., 2007), and this, combined with their high protein levels (see above) may be valuable in improving nutrition and reducing the risk of infectious diseases. Tuber conformity was varied: Ayamurasaki had the best tuber shape, size and homogeneity in the Baliem Valley, while Numfor-5 was best in Minyambouw. The best suggestion for reducing malnutrition may be that people are encouraged to plant and consume several sweetpotato cultivars with strong nutritional and agronomic traits.

Most of the introduced sweetpotato cultivars adapted well in the Baliem Valley and Minyambouw, as shown by their higher yield, particularly Pattipi, Salosa and Sawentar. Pattipi produced the highest tuber yield in the Baliem Valley, while

Sawentar produced the highest yield in Minyambouw. Bramwamrum, Helaleke, Dosak-1, Papua Pattipi, Dosak-2, Cangkuang and Ayamurasaki had higher dry matter content than other cultivars at both sites. A number of cultivars offered superior nutrient traits. Ungu, Ayamurasaki and Dosak-1, which have purple flesh colour, and also Salosa and Numfor-5, which have yellow flesh colour, had a higher protein level, while BB-00105.10, the only orange fleshed sweetpotato trialled contained the highest beta-carotene, as expected. The highest carbohydrate content was produced by the local Minyambouw cultivar (Bramwamrum), which was related to its high dry matter content.

Introduced cultivars were perceived to have the best sensory traits. Worembai was considered to have the best taste by the people of the Baliem Valley, whereas Arfak people favoured Helaleke. On the other hand, Musan, which is popular for pig diet in the Baliem Valley, was found to be fibrous and more subject to cracking than the other cultivars. It had the poorest taste according to the people of both areas.

The highland people of Papua and West Papua are recommended to grow several sweetpotato cultivars in their gardens, including high yielding forms for their daily consumption, pig feed and income, as well as cultivars that provide particular nutritional advantages.

12

Acknowledgements

The authors thank the Australian Centre for International Agricultural Research for funding, and acknowledge the input of attendees at the project's initial workshop in Manokwari, West Papua, October 2008 in conceptualising, and providing the initial plans for, this study. The International Potato Center (notably Dr Sukendra Mahalaya and Mr Koko Tjintokohadi), South Australian Research and Development Institute (Dr Colin Cargill, Project Manager) and the Indonesian partner organisations

made important input. In particular, we thank Dr Fadry Djufry, D. Tangkearing and Merlin Rumarar (Assessment Institute for Agricultural Technology Papua), Dr Wahid Rauf and Tuminem (Agriculture and Food Security Agency, West Papua), Dr Mohammed Jusuf (Research Institute for Legume and Tuber Crops, Malang), field and laboratory staff, the Directorate General of Indonesian Higher Education (DIKTI) and Dr Margaret Cargill for assistance with this article.

References

- An, V.L., Lindberg, B.E.F. and Lindberg, J.E. 2003. Effect of harvesting interval and defoliation on yield and chemical composition of leaves, stems and tubers of sweetpotato (*Ipomoea batatas* (L.) Lam) plant parts. *Field Crops Res.*, 82: 49-58.
- Association of Official Analytical Chemists (AOAC). 1990. *Official Methods of Analysis of the Association of Official Analytical Chemists*. 15th ed. Washington DC.
- Bradbury, J. H., Baines, J., Hammer, B., Anders, A. and Millar, J.S. 1984. Analysis of sweetpotato from the highlands of Papua New Guinea: Relevance to the incidence of enteritis necroticants. *J. Agric. Food Chem.*, 32: 469-473.
- Ekanayake, I.J. and Collins, W. 2004. Effect of irrigation on sweetpotato root carbohydrates and nitrogenous compounds. *Food Agric. Environ.*, 2(1): 243-248.
- Hagenimana, V., Low, J., Anyango, M., Kurz, K., Gichuki, S.T. and Kabira, J. 2001. Enhancing vitamin A intake in young children in western Kenya: Orange-fleshed sweetpotatoes and women farmers can serve as key entry points. *Food Nutr. Bull.*, 22: 376-387.
- Hill, W.A. 1984. Effect of nitrogen nutrition on quality of three important root/tuber crops. In: Hauck, R.D. (Ed.) *American Society of Agronomy, Soil Science Society of America and Crop Science Society of America, Madison, Wisconsin*. pp 627-641.
- International Potato Center (CIP). 1998. CIP sweetpotato facts. <http://www.cipotato.org/data/sweetpotato/facts/swtpfact.htm> [Accessed: 8/2/2011].
- Islam, S.M., Yoshimoto, M., Yahara, S., Okuno, S., Ishiguro, K. and Yamakawa, O. 2002. Identification and characterization of foliar polyphenolic

- composition in sweetpotato (*Ipomoea batatas* L.) genotypes. *J. Agric. Food Chem.*, 50: 3718-3722.
- Johnson, M. and Pace, D. 2010. Sweetpotato leaves: Properties and synergistic interactions that promote health and prevent disease. *Nutr. Rev.*, 68(10): 604-615.
- Jusuf, M., Setiawan, A., Peters, D., Cargill, C., Mahalaya, S., Limbongan, J. and Subandi. 2007. Memerbaiki efisiensi produksi ubi jalar-babi di Kabupaten Jayawijaya, Papua. Seminar Nasional dan Ekspose Percepatan Inovasi Teknologi Pertanian Spesifik Lokasi, Jayapura 5-6 Juni 2007 (in Indonesian language).
- Kano, M.T., Takayagi, K., Harada, K., Makino, F. and Ishikawa. 2005. Antioxidative activity of anthocyanins from purple sweetpotato, *Ipomoea batatas* cultivar Ayamurasaki. *Biosci., Biotechnol., Biochem.*, 69(5): 979-988.
- Kumar, B.M. 2006. Agroforestry: the new old paradigm for Asian food security. *J. Trop. Agric.*, 44: 1-14.
- Lila, M.A. 2004. Anthocyanins and human health: An in vitro investigative approach. *J. Biomed. Biotech.*, 5: 306-313.
- Oboh, S., Ologhobo, A. and Tewe, O. 1989. Some aspects of the biochemistry and nutritional value of the sweetpotato (*Ipomoea batatas*). *Food Chem.*, 31:9-18.
- Ofori, G., Oduro, I., Ellis, W.O. and Dapaah, K.H. 2009. Assessment of vitamin A content and sensory attributes of new sweetpotato (*Ipomoea batatas*) genotypes in Ghana. *Afr. J. Food Sci.*, 3(7): 184-192.
- Peters, J. 2001. Local human-sweetpotato-pig systems characterization and research in Irian Jaya, Indonesia. With limited reference to Papua New Guinea. A secondary literature review. Unpublished report of ACIAR AH/1998/054 Project. International Potato Center (CIP) Bogor.
- Rahman, S.M.M., Wheatley, C. and Rakshit, K.S. 2003. Selection of sweetpotato variety for high starch extraction. *Inter. J. Food Prop.*, 6(3): 419-430.
- Rodriguez-Amaya, D. and Kimura, M. 2004. HarvestPlus Handbook for Carotenoid Analysis. HarvestPlus technical Monograph 2. International Food Policy Research Institute and International Center for Tropical Agriculture, Washington DC, USA and Cali, Colombia.
- Rainard, J. 1969. Notes on sweetpotato research in West New Guinea (West Irian). In: Tai, E.A., Charles, W.B., Iton, E.F., Hayness, P.H., and Leslie, K.A (eds), Proceedings of the international symposium on tropical root crops. Vol. I. University of the West Indies, Trinidad. pp 88-111.
- Schneider, J., Widyastuti, C.A., and Djazuli, M. 1993. Sweetpotato in the Baliem Valley area, Irian Jaya. In: A report on collection and study of sweetpotato germplasm, April-May 1993. International Potato Center (CIP), ESEAP-Region, Bogor, Indonesia.
- Sillitoe, P. 1998. It's all in the mound: fertility management under stationary shifting cultivation in the Papua New Guinea Highlands. *Mountain Res. Dev.*, 18(2): 123-134.
- Soenarto and Rumawas, F. 1997. An agro-ecological analysis of *wen-tinak*, a sustainable sweetpotato wetland production system in the Baliem Valley, Irian Jaya, Indonesia. *Sci. New Guinea*, 23(2): 55-66.
- Soplanit, A., Wanaer, D and Tirajoh, S. 2005. Pengkajian pemupukan organik dan jarak tanam pada tanaman ubi jalar di dataran tinggi Jayawijaya. In: Prosiding Seminar Nasional Optimalisasi Teknologi Kreatif dan Peran Stakeholder dalam Percepatan Adopsi Inovasi Teknologi Pertanian. Denpasar, 28 September 2005 (in Indonesian language).
- Takahata, Y., Noda, T. and Nagata, T. 1993. Varietal differences in chemical composition of the sweetpotato storage root. *Acta Hort.*, 343: 77-80.
- Teow, C.C., Truong, V.D., Mc. Feeters, R.F., Thompson, R.L., Pecota, K.V. and Yencho, G.C. 2007. Antioxidant activities, phenolic and β -carotene contents of sweetpotato genotypes with varying flesh colours. *Food Chem.*, 103: 829-838.
- Woolfe, J.A. 1992. Sweetpotato: An untapped food resource. Cambridge University Press; Cambridge, United Kingdom.
- Yamakawa, O. and Yoshimoto, M. 2002. Sweetpotato as food material with physiological functions. *Acta Hort.* (ISHS), 583: 179-85.

Yield trial and sensory evaluation of sweetpotato cultivars in Highland Papua and West Papua Indonesia

ORIGINALITY REPORT

13%

SIMILARITY INDEX

9%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Wageningen University 2%
Student Paper
- 2** researchspace.ukzn.ac.za 1%
Internet Source
- 3** Babajide Olusegun Omotobora, Patrick Olusanmi Adebola, David Mxolisi Modise, Sunette Marlize Laurie, Abe Shegro Gerrano. "Greenhouse and Field Evaluation of Selected Sweetpotato (*Ipomoea batatas* (L.) LAM) Accessions for Drought Tolerance in South Africa", American Journal of Plant Sciences, 2014 1%
Publication
- 4** M Hayati, Sabaruddin, Efendi, A Anhar. "Chemical and sensory analysis of several sweet potatoes (L.) clones", IOP Conference Series: Earth and Environmental Science, 2019 1%
Publication
- 5** www.mdpi.com 1%
Internet Source

6	repository.up.ac.za Internet Source	1 %
7	researchonline.jcu.edu.au Internet Source	1 %
8	www.ijcrar.com Internet Source	1 %
9	Sochinwechi Nwosisi, Dilip Nandwani, Dafeng Hui, Ramasamy Ravi. "Sensory Evaluation of Organic Sweetpotato Cultivars", International Journal of Vegetable Science, 2017 Publication	<1 %
10	core.ac.uk Internet Source	<1 %
11	sevgiligiyim.com Internet Source	<1 %
12	www.regional.org.au Internet Source	<1 %
13	M. R. Bakker. "Root distribution of <i>Fagus sylvatica</i> in a chronosequence in western France", Journal of Forest Research, 06/2008 Publication	<1 %
14	Tekalign, T.. "Growth and productivity of potato as influenced by cultivar and reproductive growth", Scientia Horticulturae, 20050530 Publication	<1 %

15	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
16	docksci.com Internet Source	<1 %
17	repository.out.ac.tz Internet Source	<1 %
18	www.crawfordfund.org Internet Source	<1 %
19	Submitted to Rhodes University Student Paper	<1 %
20	www.science.gov Internet Source	<1 %
21	M. Jusuf Yakub, D. Peters. "ADAPTATION AND YIELD STABILITY OF SWEETPOTATO CLONES FOR HUMAN AND PIG FOOD IN MOUNTAINOUS AREAS OF JAYAWIJAYA, IRIAN JAYA, INDONESIA", Acta Horticulturae, 2005 Publication	<1 %
22	journals.ametsoc.org Internet Source	<1 %
23	www.eseap.cipotato.org Internet Source	<1 %
24	www.scribd.com Internet Source	<1 %

25 V. HAGENIMANA. "OIL CONTENT IN FRIED PROCESSED SWEETPOTATO PRODUCTS", Journal of Food Processing and Preservation, 7/1998
Publication <1 %

26 ageconsearch.umn.edu
Internet Source <1 %

27 www.sua.ac.tz
Internet Source <1 %

28 J. C. Van Dam, R. Singh, J. J. E. Bessembinder, P. A. Leffelaar, W. G. M. Bastiaanssen, R. K. Jhorar, J. G. Kroes, P. Droogers. "Assessing Options to Increase Water Productivity in Irrigated River Basins Using Remote Sensing and Modelling Tools", International Journal of Water Resources Development, 2006
Publication <1 %

29 Thybo, A.. "Low-field NMR relaxation and NMR-imaging as tools in differentiation between potato sample and determination of dry matter content in potatoes", LWT - Food Science and Technology, 200305
Publication <1 %

30 article.sciencepublishinggroup.com
Internet Source <1 %

31 dspace.knust.edu.gh
Internet Source <1 %

32

erl.ucc.edu.gh:8080

Internet Source

<1 %

33

pdfs.semanticscholar.org

Internet Source

<1 %

34

www.zef.de

Internet Source

<1 %

35

Fekadu Gurmu, Shimelis Hussein, Mark Laing. "Evaluation of candidate orange-fleshed sweetpotato clones for nutritional traits", *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science*, 2017

Publication

<1 %

36

Ruth W. Mukhongo, John B. Tumuhairwe, Peter Ebanyat, AbdelAziz H. AbdelGadir, Moses Thuita, Cargele Masso. "Combined Application of Biofertilizers and Inorganic Nutrients Improves Sweet Potato Yields", *Frontiers in Plant Science*, 2017

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Yield trial and sensory evaluation of sweetpotato cultivars in Highland Papua and West Papua Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
