

# In vivo Efficacy of *Biophytum petersianum* on *Haemonchus contortus* in Goats

*by* Priyo Sambodo

---

**Submission date:** 16-Apr-2020 02:58PM (UTC+0800)

**Submission ID:** 1298980669

**File name:** AAVS\_8\_3\_238-244.pdf (506.45K)

**Word count:** 4776

**Character count:** 25130



## *In vivo* Efficacy of *Biophytum petersianum* on *Haemonchus contortus* in Goats

PRIYO SAMBODO<sup>1</sup>, JOKO PRASTOWO<sup>2</sup>, KURNIASIH KURNIASIH<sup>3</sup>, WIDA WAHIDAH MUBAROKAH<sup>4</sup>, SOEDARMANTO INDARJULIANTO<sup>5\*</sup>

<sup>1</sup>Department of Animal Science Papua University, Jalan Gunung Salju, Amban, Manokwari Papua Barat, 98314, Indonesia; <sup>2</sup>Department of Parasitology, Faculty of Veterinary, Universitas Gadjah Mada, Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281, Indonesia; <sup>3</sup>Department of Pathology, Faculty of Veterinary, Universitas Gadjah Mada, Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281, Indonesia; <sup>4</sup>Department of Animal Health, Agriculture Extension College, The Polytechnique of Agricultural Development Yogyakarta Magelang, Jalan Magelang-Kopeng Km 7 Purwosari, Tegalsrejo, Magelang, Jawa Tengah, 56192, Indonesia; <sup>5</sup>Department of Internal Medicine, Faculty of Veterinary, Universitas Gadjah Mada, Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281, Indonesia.

**Abstract** | Haemonchosis is a common and severe disease caused by the infection of worms *Haemonchus* sp. Kebar grass (*Biophytum petersianum*) known as a plant from Papua Indonesia that contains tannin compounds and has potential anthelmintic activity. The present study was carried out to determine the influence of Kebar grass infusion on *H. contortus* under *in vivo* conditions. A total of 15 female goats at 6-8 months old were divided into 5 groups, and each consisted of 3 goats. All groups were infected 1000 infective larvae orally every week for 4 weeks. Three groups were given Kebar grass infusion a day at a dose of 2 mg/mL, 4 mg/mL and 6 mg/mL respectively for 7 days at the 6th week. Group 4 was given Albendazole at a dose of 3.8 mg/kg body weight, and group 5 was not given treatment. Clinical examinations, hematological and EPG were carried out every week. All goats were autopsied and analyzed for pathology at the seventh week. Quantitative data to perform statistical analysis, as well as the results of physical examination and histopathology were analyzed descriptively. Body weight measurements *in vivo* as the mean, Body Condition Scores (BCS), and FAMACHA showed an increase. RBC, Hb, PCV and total protein values increased. The FECR value of the treatment group was higher than the negative control group. During the autopsy, both carcass and visceral organs were colourless, slight of subcutaneous fat, and small intestine haemorrhage and ascites in the abdominal cavity. Many nodules with diameters between 1.5-2.0 mm were observed in the abomasal mucosa. Histopathological changes in the control group were infiltration of inflammatory cells, congestion and the presence of worm pieces. Based on results obtained in this line of research, it is therefore concluded that Kebar grass infusion is anthelmintic against *H. contortus* under *in vivo* conditions.

**Keywords** | Anthelmintic, *Haemonchus contortus*, Goats, Kebar grass

**Received** | October 07, 2019; **Accepted** | February 17, 2020; **Published** | March 05, 2020

\***Correspondence** | Sudarmanto Indarjulianto, Department of Internal Medicine, Faculty of Veterinary, Universitas Gadjah Mada, Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281, Indonesia; **Email:** indarjulianto@ugm.ac.id

**Citation** | Sambodo P, Prastowo J, Kurniasih K, Mubarakah WW, Indarjulianto S (2020). *In vivo* efficacy of *Biophytum petersianum* on *Haemonchus contortus* in goats. Adv. Anim. Vet. Sci. 8(3): 238-244.

**DOI** | <http://dx.doi.org/10.17582/journal.aavs/2020/8.3.238.244>

**ISSN (Online)** | 2307-8316; **ISSN (Print)** | 2309-3331

**Copyright** © 2020 Sambodo *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Haemonchosis is one of the major parasitic diseases in goats in Indonesia. The biggest economic losses due to haemonchosis occur in sub-tropical and tropical regions

which are mainly due to mortality, decreased production, stunted growth and low body weight (Mengist *et al.*, 2014). In fact, West Papua could be a potential development area for goat livestock. Based on data from the Department of Animal Husbandry of West Papua Province (2016), the



number of goats in West Papua Province has continued to increase by more than 77% within 4 years. One obstacle in developing the livestock business is the high incidence of helminthiasis. With regard to this issue, Purwaningsih et al. (2017), stated that the incidence of gastrointestinal worm infestations in etawa-bred goats that were kept semi-intensively in Amban District, Manokwari was 100% (n: 32). At present, resistance to nematode populations has been detected in all livestock species, such as sheep, goats, cattle and horses (Kaplan, 2004). From actual cases appear, Kebar grass (*Biophytum petersianum* Klotzsch) developed as one of Indonesia's endemic plants used as medicinal plant. The plant tannin content is 1.1% (Santoso et al., 2006) and condensed tannins are 0.28% (Sambodo, 2018). As well, Widayati and Nurhayati (2013), reported that at 60% dose of *Biophytum petersianum* Klotzsch is able to eradicate adult worm called *Ascaridia galli* in 7 hours which is used as sample. On the other hand, 60% dose infusion of *Biophytum petersianum* works independently to also execute those adults worms in 7.5 hours under *in vivo* condition. From such observations, it can be argued that the Kebar grass has potential as an anthelmintic of *H. contortus* worms under *in vivo* conditions.

## MATERIAL AND METHOD

### ETHICAL CLEARANCE

The entire research processes have received ethical eligibility from the Ethics Commission of the Gadjah Mada University in Yogyakarta (Certificate No.: 00116/04/LPPT/IX/2017).

### COLLECTION AND BIOPHYTUM PETERSIANUM INFUSION

*Biophytum petersianum* was obtained from the Central Kebar District (located at 133°03'25.8" East Longitude and 00°48'31.1" South Latitude with an altitude of 590 m asl) Tambrauw Regency, West Papua Province. First of all, some Kebar grass is aired, thus the Kebar grass is dried and stored until it is used for the extraction process (Sambodo et al., 2012).

The kebar grass is dried and finely chopped. A total of 1 gram (1%), 2 gram (2%), 4 gram (4%), 6 gram (6%), 8 gram (8%) and 10 gram (10%) dried Kebar grass were put into the Erlenmeyer and soaked in 100 mL aqua destillata. Erlenmeyer flask was incubated in an oven at 90 °C for 15 minutes. Furthermore the solution is filtered using filter paper and stored in a refrigerator for 24 hours (Daryatmo et al., 2010).

### TREATMENT ON EXPERIMENTAL GOATS

The treatment was carried out according to Eguale et al. (2007). Experimental animals were 15 female goats aged 6-8 months divided into 5 groups. Four weeks after adaptation, all groups were infected with 1000 oral infective

larvae every week for 4 weeks. At an average EPG value of 3000 EPG, all groups were given treatment for 7 days as follows group 1 (2 mg/mL), group 2 (4 mg/mL), group 3 (6 mg/mL), group 4/positive control (3.8 mg/kg BW) and group 5/negative control (not treated). All research parameter checks were carried out on the 0th and 7th days.

### CLINICAL EXAMINATION

Clinical examination included weighing and physical examination in forms of eye conjunctiva (FAMACHA) and BCS (Eguale et al., 2007; Attindehou et al., 2012).

### HEMATOLOGY

Blood collection was carried out on all goats through the jugular vein of 2 ml/ goat. The parameters analyzed included RBC, Hb, PCV and total protein (Eguale et al., 2007; Qamar and Maqbool, 2012).

### FECAL EGG COUNT REDUCTION (FECR)

Fecal sample collection was performed directly from the rectum. The EPG calculation was carried out using the McMaster technique according to Coles et al. (1992). The percentage decrease in the number of worm eggs per gram of faeces or FECR was calculated by the following guidelines:

$$\text{FECR (\%)} = \frac{\text{EPG pre-treatment} - \text{EPG post-treatment}}{\text{EPG pre-treatment}} \times 100$$

### POSTMORTEM EXAMINATION

The necropsy technique was carried out mainly according to Somvanshi and Rao (2009), with several modifications. After physical examination, the goats are slaughtered by cutting the four neck veins. Move forward to perform the dissection to the dead goats. Examination of the carcass, visceral organ and intestine as well as the presence or absence of fluid in the abdominal cavity. Macroscopic examination of mucosal abomasum and tissue collection as histological preparations were made according to Muntiha (2001).

### DATA ANALYSIS

The study adopted quantitative data to perform statistical analysis. The findings were presented as mean ± standard deviation. The difference from the effect of different treatments was analyzed by analysis of variance with SPSS 23 (IBM Corp). The correlation between physical examination and histopathological data were presented descriptively.

## RESULTS

### CLINICAL EXAMINATION

In the current project, the mean body weight of goats increased post-treatment compared to pre-treatment (Table 1), although paired t-tests did not produce a




significant difference ( $P>0.01$ ) and the value of BCS has increased, yet it is still in the low range. The value of FAMACHA after treatment increased compared to pre-treatment, where treatment on group 3 experienced the greatest increase (Figure 1).

**Table 1:** Average weight of the treatment group.

Group	Body weight	
	Pre-treatment	Post-treatment
Group 1 (2 mg/mL)	12.67±1.15 <sup>a</sup>	13.50±1.00 <sup>a</sup>
Group 2 (4 mg/mL)	14.00±1.32 <sup>a</sup>	14.50±2.50 <sup>a</sup>
Group 3 (6 mg/mL)	13.00±1.00 <sup>a</sup>	13.67±1.53 <sup>a</sup>
Average	13.22±1.18 <sup>a</sup>	13.89±1.62 <sup>a</sup>

The same superscript in the same line are not significantly different ( $P>0.005$ ).



**Figure 1:** FAMACHA value of the treatment group. (a) Pre-treatment (value 5) (b) post-treatment (value 3).

**Table 2:** Mean haematological values of pre-treatment and post-treatment.

Group	Parameters	Pre-treatment	Post-treatment
Control + (Albendazole 3.8 mg/kg BW)	RBC ( $\mu$ L)	2.23±0.38 <sup>a</sup>	2.25±0.31 <sup>a</sup>
	Hb (dL)	9.00±0.82 <sup>a</sup>	9.30±0.60 <sup>a</sup>
	PCV (%)	25.00±4.21 <sup>a</sup>	26.00±3.91 <sup>a</sup>
	Serum protein (g/dl)	6.60±0.32 <sup>a</sup>	6.70±0.44 <sup>a</sup>
Group 1 (2 mg/mL)	RBC ( $\mu$ L)	1.92±0.84 <sup>a</sup>	2.11±0.45 <sup>a</sup>
	Hb (dL)	8.00±2.53 <sup>a</sup>	7.30±2.61 <sup>a</sup>
	PCV (%)	20.00±11.45 <sup>a</sup>	18.95±9.05 <sup>a</sup>
	Serum protein (g/dl)	6.00±0.45 <sup>a</sup>	6.40±0.48 <sup>a</sup>
Group 2 (4 mg/mL)	RBC ( $\mu$ L)	2.27±0.75 <sup>a</sup>	2.75±0.88 <sup>a</sup>
	Hb (dL)	8.00±1.58 <sup>a</sup>	9.00±1.56 <sup>a</sup>
	PCV (%)	25.00±8.26 <sup>a</sup>	30.00±10.72 <sup>a</sup>
	Serum protein (g/dl)	7.00±0.11 <sup>a</sup>	7.00±0.82 <sup>a</sup>
Group 3 (6 mg/mL)	RBC ( $\mu$ L)	1.91±0.11 <sup>a</sup>	2.67±0.49 <sup>a</sup>
	Hb (dL)	7.00±3.27 <sup>a</sup>	9.00±1.17 <sup>a</sup>
	PCV (%)	20.00±10.91 <sup>a</sup>	30.00±4.97 <sup>a</sup>
	Serum protein (g/dl)	3.00±0.51 <sup>a</sup>	7.00±0.15 <sup>a</sup>
Control - (saline water 0,62%)	RBC ( $\mu$ L)	2.33±0.54 <sup>a</sup>	2.01±0.49 <sup>a</sup>
	Hb (dL)	8.90±1.25 <sup>a</sup>	7.90±1.21 <sup>a</sup>
	PCV (%)	23.30±3.79 <sup>a</sup>	19.80±3.56 <sup>a</sup>
	Serum protein (g/dl)	6.90±0.93 <sup>a</sup>	5.00±2.03 <sup>a</sup>

The same superscript in the same line are not significantly different ( $P>0.005$ ).

**HEMATOLOGY**

RBC, Hb, PCV and total protein values post-treatment with Kebar grass infusion showed an increase compared to pre-treatment, where for all parameters, treatment group 3 experienced the greatest increase (Table 2).

**FECR**

At this stage, the average EPG for the 3 treatment groups was 2955 EPG. The average EPG pre and post treatment as well as the results of FECR calculations in each treatment group can be illustrated in Table 3.

**Table 3:** Average EPG and FECR treatment groups.

Groups	EPG		FECR (%)
	Pre-treatment	Post-treatment	
Control + (Albendazole 3.8 mg/kg BW)	1583±575.18	0±0.00	100.00
Group 1 (2 mg/mL)	2217±125.83 <sup>a</sup>	0±0.00 <sup>b</sup>	100.00
Group 2 (4 mg/mL)	3650±2995.41 <sup>a</sup>	0±0.00 <sup>b</sup>	100.00
Group 3 (6 mg/mL)	3000±2133.66 <sup>a</sup>	17±28.87 <sup>b</sup>	99.44
Control - (saline water 0.62%)	1900±888.82	1450±390.51	23.68

a,b means with different superscripts in the line are significantly different ( $P<0.005$ ).



**Figure 2:** Mucosa abomasum control and treatment group. (a) positive control; (b) treatment 1, nodule (arrow); (c) treatment 2; (d) treatment 3; (e) negative controls, nodules (arrow).

**POSTMORTEM EXAMINATION**

**ANATOMICAL PATHOLOGY**


In the present study, the carcass condition between the negative control group and the treatment group was not relatively different, the two carcasses were colorless and



subcutaneous fat was highlighted in small amount, the visceral organs of the negative control group appeared paler and the small intestine in both groups showed hemorrhages. There was ascites fluid in the abdominal cavity, although the amount was more in the negative control group. In the abomasal mucosa the negative control group and treatment on group 1 had many nodules with a diameter between 1.5-2.0 mm (Figure 2).

### HISTOLOGY

Histopathological examination revealed abnormal conditions in both the control and treatment groups (Figures 3). In abomasum tissue, it was found inflammation infiltration, congestion, larvae 4 and adult worms.



**Figure 3:** Histopathology of the abomasum control group. (a) positive control, larvae 4 in the submucosa (arrow) (40x); (b) positive control, congestion of blood vessels in the submucosa (20x); (c) group 1, infiltration of inflammatory

cells in the sub mucosa (arrow) (40x); (d) group 1, edema in the sub mucosa (4x); (e) group 2, infiltration of inflammatory cells in the sub mucosa (arrow) (20x); (f) group 2, inflammatory cell infiltration (black arrow) and edema in the sub mucosa (green arrow) (40x); (g) group 3, infiltration of inflammatory cells in the sub mucosa (arrow) (40x); (h) group 3, congestion of sub mucosal blood vessels (40x); (i) negative control, cross section of adult worms in the mucosa (arrow) (20x); (j) negative control, infiltration of inflammatory cells in the sub mucosa (arrow) (40x).

### DISCUSSION

#### CLINICAL EXAMINATION

Aside from the experimental goats used at a young age of 6-8 months, the low body weight and BCS in this study were directly proportional to the low RBC value and total serum protein due to helminth infections. Regarding this issue, Eguale et al. (2007) reported, goats kept in animal husbandry and given dry feed; lose blood plasma and protein due to parasites which is would prevent goats to gaining weight.

The level of anemia can be roughly evaluated by looking at the color of the conjunctiva where the blood capillaries are on the surface, thus the color of the tissue reflects the color of the blood. In general, the value of FAMACHA in this study is in the normal range (3), that in such formulated category, the goats do not need to be given anthelmintics even though Vatta et al. (2001), reported that the goats in the FAMACHA 3, 4 and 5 categories were still mainly considered to be given anthelmintics, because the conjunctival color range is smaller in goats than sheep, and as a result, the application process of the FAMACHA system slightly can be difficult to apply. In addition, the sensitivity only reaches 67-69%.

#### HEMATOLOGY

An increase in each parameter after treatment was remarkably considered to be due to the reduction or absence of adult worms in the treatment group, evidenced by the results of the necropsy abomasum of the treatment group that no longer found adult worms. The RBC value and total serum protein are lower than the reference according to Rasedee (1981). This decrease in RBC may be caused by the amount of blood losses during an infection. Each worm can suck 0.05 ml of blood per day (Urquhart et al., 1996). The decrease in total serum protein may be due to blood thinning due to abomasal bleeding caused by larval invasion (Qamar and Maqbool, 2012).

#### FECR

During the treatment, no poisoning effect was found in all experimental animals. In addition, at the time of necropsy,

no adult worms were found in all abomasum from the positive treatment and control group. EPG values before and after treatment decreased significantly ( $P = 0.00$ ). The results of this study are in accordance with previous studies. Iqbal *et al.* (2004), stated that a dose of 3 g / kg BW of water extract of *Artemisia brevifolia* was able to reduce the EPG value of sheep GI nematodes by 67.2% and was added by Tariq *et al.* (2009), that a dose of 2 gm / kg BW of water extract *Artemisia absinthium* was able to reduce the EPG value of sheep GI nematodes by 80.49%. *Artemisia* plants are rich in proanthocyanidins (condensed tannins) (Lutgen, 2018). Athanasiadou *et al.* (2001), reported that FEC and worms in the small intestine would decrease in number in sheep fed with pellets containing condensed tannins from Quebracho tree extracts. Sheep fed with Sainfoin straw containing dry condensed tannins decreased FEC (Paolini *et al.*, 2003) and those fed *Acacia karoo* dried leaves decreased both FEC and the number of worms (Kahiya *et al.*, 2003).

In this study, it was believed that the effects of condensed tannins occurred directly. The condensed tannins in the Kebar grass extract directly inhibit the hatching of worm eggs, where the EPG value drops significantly. In addition, condensed tannins also binds to protein, where under *in vitro* test there is a decrease in the number of protein bands and body surface damage in adult worms, and therefore, they become inactive and finally die.

#### POSTMORTEM EXAMINATION ANATOMICAL PATHOLOGY

These results are mainly caused by *H. contortus* infestation. *Haemonchus contortus* suck blood which causes blood loss around the tissue make it looks colorless and resulting in decreased body weight. On the other hand, blood loss will drop blood albumin levels and cause fluid accumulation in the abdominal cavity. Both adult and fourth stage larvae suck blood and in addition, migration of adult and larvae cause haemorrhages into the abomasum.

Saminathan *et al.* (2015) suggested that in sheep infected with *H. contortus* has pale carcasses, little subcutaneous fat, yellowish fluid in the abdominal cavity, pale internal organs and small intestine congested, watery and petechiae hemorrhage and added by Estuningsih *et al.* (1996), that the surface of the abomasum infected with *H. contortus* showed changes in the form of lesions, inflammation due to worm bites. Edema of the folds of the omasum, hemorrhage of petechiae and the formation of nodules in the infected abomasum may be caused by the activity of the worm while sucking blood (Soulsby, 1982).

#### HISTOPATOLOGY

Attachment of the parasite with the abomasum mucosal causes histopathological changes and infection.

Histopathological changes caused by *H. contortus* infection, such as the occurrence of a strong inflammatory response in the infected area characterized by cellular infiltration in the form of fibroblasts, macrophages, lymphocytes, plasma cells, mast cells, neutrophils and eosinophils (Balic *et al.*, 2002; Ortolani *et al.*, 2013; Saminathan *et al.*, 2015).

Although many behaviors could be considered as abnormal in the treatment group, it was also believed that the major cause was due to the induction of previous haemonchosis. Although therapy has been conducted and no adult worms were found on the surface of the abomasum, the condition of goats tried to influence the healing process. The ability of the host to develop immunity and the level of resistance against GI nematodes depends on many factors, such as type, age, number of worms and nutrition (Stear and Murray, 1994; Hoste *et al.*, 2005; Rowe *et al.*, 2008). Pathology caused by parasites and host reactions to infecting parasites depend on the organ being attacked, depth of penetration, location and number of parasites (Taraschewski, 2000; Esmailnejad *et al.*, 2012; de Oliveira *et al.*, 2013) and added by Hoste *et al.* (2008), compared to sheep, goats develop fewer immune responses against nematodes, which in turn might affect the state of pathology and the ability to control infection.

Research on this issue, the presence of larvae (L4) in the abomasum tissue of the group given Albendazole, may be due to the low dose of the drug that contacts the parasite when it enters the body of the host. The metabolism of an anthelmintic in a host affects the anthelmintic concentration in the parasite (Lifschitz *et al.*, 2017).

#### CONCLUSION

Based on clinical examination, hematological values, FECR values and postmortem examination *in vivo*, Kebar grass infusion is anthelmintic against *H. contortus*.

#### ACKNOWLEDGMENTS

This study was supported by the Ministry of Research, Technology and Higher Education of the Republic of Indonesian through Domestic Graduate Scholarship (BPP-DN) program, grant number 1265.35/E4.4/2015.

#### CONFLICT OF INTEREST

The authors declare that they have no competing interest.

#### AUTHORS CONTRIBUTION

PS, JP, KK and SI the researchers. PS, JP, KK and SI devel-



oped the research concepts. PS and JP prepared *H. contortus*, PS, WWM and SI managed and examined goats during *in vivo* experiments and interpreted the results; PS and KK made histopathological preparations and interpreted the results; PS and WWM collected data and literature and prepared manuscripts. JP, KK and SI guided and corrected and complete the manuscript.

## REFERENCES

- Athanasiadou SL, Kyriazakis I, Jackson F, Coop RL (2001). Direct anthelmintic effects of condensed tannins towards different gastrointestinal nematodes of sheep: *in vitro* and *in vivo* studies. *Vet. Parasitol.* 99(3): 205-219. [https://doi.org/10.1016/S0304-4017\(01\)00467-8](https://doi.org/10.1016/S0304-4017(01)00467-8)
- Attindehou S, Salifou S, Biaou CF, Gbati OB, Adamou-N'diaye M, Pangui LJ (2012). Epidemiology of *Haemonchosis* in sheep and goats in Benin. *J. Parasitol. Vect. Biol.* 4(2): 20-24.
- Balic A, Bowles VM, Meeusen ENT (2002). Mechanisms of immunity to *Haemonchus contortus* infection in sheep. *Parasite Immunol.*, 24(1): 39-46. <https://doi.org/10.1046/j.0141-9838.2001.00432.x>
- Coles GC, Bauer C, Borgsteede FHM, Geerts S, Klei TR, Taylor MA, Waller PJ (1992). World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) methods for the detection of anthelmintic resistance in nematodes of veterinary importance. *Vet. Parasitol.* 44(1-2): 35-44. [https://doi.org/10.1016/0304-4017\(92\)90141-U](https://doi.org/10.1016/0304-4017(92)90141-U)
- Daryatmo J, Hartadi H, Orskov ER, Adiwimarta K, Nurcahyo W (2010). *In vitro* screening of various forages for anthelmintic activity on *Haemonchus contortus* eggs. *Adv. Anim. Biosci.* 1(1): 113-113. <https://doi.org/10.1017/S2040470010002566>
- de Oliveira RB, Senger MR, Vasques LM, Gasparotto J, dos Santos JPA, de Bittencourt Pasquali MA, Moreira JCF, Silva FP, Gelain DP (2013). *Schistosoma mansoni* infection causes oxidative stress and alters Receptor for Advanced Glycation Endproduct (RAGE) and tau levels in multiple organs in mice. *Int. J. Parasitol.* 43(5): 371-379. <https://doi.org/10.1016/j.ijpara.2012.12.006>
- Department of Animal Husbandry of West Papua Province (2016). West Papua Province Livestock Statistics Book 2011-2015. Manokwari, West Papua.
- Eguale T, Tilahun G, Debella A, Feleke A, Makonnen E (2007). *In vitro* and *in vivo* anthelmintic activity of crude extracts of *Coriandrum sativum* againsts *Haemonchus contortus*. *J. Ethnopharmacol.* 110(3): 428-433. <https://doi.org/10.1016/j.jep.2006.10.003>
- Esmailnejad B, Tavassoli M, Asri-Rezaei S, Dalir-Naghadeh B (2012). Evaluation of antioxidant status and oxidative stress in sheep naturally infected with *Babesia ovis*. *Vet. Parasitol.* 185(2): 124-130. <https://doi.org/10.1016/j.vetpar.2011.10.001>
- Estuningsih S, Retnani EB, Esfandiari A (1996). The pathological changes of male goat infected by *Haemonchus contortus* (Rudolphi, 1803). *Med. Vet.* 3(2): 39-52.
- Hoste H, Torres-Acosta JF, Paolini V, Aguilar-Caballero A, Etter E, Lefrileux Y, Chartier C, Broqua C (2005). Interactions between nutrition and gastrointestinal infections with parasitic nematodes in goats. *Small Ruminant Res.* 60(1-2): 141-151. [- \[smallrumres.2005.06.008\]\(https://doi.org/10.1016/j.jep.2004.03.046\)
  - Hoste H, Torres-Acosta JF, Aguilar-Caballero A \(2008\). Nutrition-parasite interactions in goats: is immunoregulation involved in the control of gastrointestinal nematodes? \*Parasite Immunol.\* 30\(2\): 79-88.
  - Iqbal Z, Lateef M, Ashraf M, Jabbar A \(2004\). Anthelmintic activity of \*Artemisia brevifolia\* in sheep. \*J. Ethnopharmacol.\* 93\(2-3\): 265-268. <https://doi.org/10.1016/j.jep.2004.03.046>
  - Kahiya C, Mukairwa S, Thamsborg, SM \(2003\). Effects of \*Acacia nilotica\* and \*Acacia karoo\* diets on \*Haemonchus contortus\* infection in goats. \*Vet. Parasitol.\* 115\(3\): 265-274. \[https://doi.org/10.1016/S0304-4017\\(03\\)00213-9\]\(https://doi.org/10.1016/S0304-4017\(03\)00213-9\)
  - Kaplan, R.M. 2004. Drug resistance in nematodes of veterinary importance. A status reports. \*Trends Parasitol.\* 20\(10\): 477-481. <https://doi.org/10.1016/j.pt.2004.08.001>
  - Lifschitz A, Lanusse C, Alvarez L \(2017\). Host pharmacokinetics and drug accumulation of anthelmintics within target helminth parasites of ruminants. \*N. Z. Vet. J.\* 65\(4\): 176-184. <https://doi.org/10.1080/00480169.2017.1317222>
  - Lutgen P \(2018\). Tannins in artemisia: the hidden treasure of prophylaxis. \*Pharm. Pharmacol. Int. J.\* 6\(3\): 176-181. <https://doi.org/10.15406/ppij.2018.06.00173>
  - Mengist Z, Abebe N, Gugsa G, Kumar N \(2014\). Assessment of small ruminant \*Haemonchosis\* and its associated risk factors in and around Finoteselam, Ethiopia. \*J. Agric. Vet. Sci.\* 7\(12\): 36-41. <https://doi.org/10.9790/2380-071223641>
  - Muntiha M \(2001\). Histopathology preparation techniques from animal tissues with Hematoxylin and Eosin \(H and E\) staining. Non-researcher functional technical meeting. \*Agric. Res. Dev. Center Minist. Agric.\* pp. 156-163.
  - Ortolani EL, Leal MLR, Minervino AHH, Aires AR, Coop RL, Jackson F, Suttle NF \(2013\). Effects of parasitism on cellular immune response in sheep experimentally infected with \*Haemonchus contortus\*. \*Vet. Parasitol.\* 196 \(1-2\): 230-234. <https://doi.org/10.1016/j.vetpar.2013.02.014>
  - Paolini V, Dorchie P, Hoste H \(2003\). Effects of sainfoin hay on gastrointestinal infection with nematodes in goats. \*Vet. Record.\* 152\(19\): 600-601. <https://doi.org/10.1136/vr.152.19.600-b>
  - Purwaningsih, Noviyanti, Sambodo P \(2017\). Gastrointestinal helminth infestation on ettawa crossbreed goat in Amban Subdistrict Manokwari Barat District Manokwari Regency West Papua Province. \*J. Il. Pet. Terpadu.\* 5\(1\): 8-12. <https://doi.org/10.23960/jipt.v5i1.p8-12>
  - Qamar MF, Maqbool A \(2012\). Biochemical studies and serodiagnosis of \*haemonchosis\* in sheep and goats. \*J. Anim. Plant Sci.\* 22\(1\): 32-38.
  - Rasdee A \(1981\). Blood parameters of kambing katjang-jumnapari crossbred goats in Malaysia. \*Pertanika.\* 4\(1\): 87-90.
  - Rowe A, McMaster K, Emery D, Sangster N \(2008\). \*Haemonchus contortus\* infection in sheep: parasite fecundity correlates with worm size and host lymphocyte counts. \*Vet. Parasitol.\* 153\(3-4\): 285-293. <https://doi.org/10.1016/j.vetpar.2008.01.040>
  - Santoso B, Kilmaskossu A, Sambodo P \(2006\). Effects of saponin from \*Biophytum petersianum\* Klotzsch on ruminal fermentation, microbial protein synthesis and nitrogen utilization in goats. \*Anim. Feed Sci. Tech.\* 137 \(1-2\): 58-68. <https://doi.org/10.1016/j.anifeeds.2006.10.005>
  - Sambodo P, Prastowo P, Indarjulianto S \(2012\). Larvicidal](https://doi.org/10.1016/j.</a></li>
</ul>
</div>
<div data-bbox=)

activity of the rumpup kebar (*Biophytum petersianum* Klotzsch) ethanol extract againsts *Aedes aegypti*. J. Il Pet. 7(1): 8-10.

- 15 • Sambodo P, Prastowo J, Kurniasih K, Indarjulianto S (2018). *In vitro* potential anthelmintic activity of *Biophytum petersianum* on *Haemonchus contortus*. Vet. World. 11(1): 1-4. <https://doi.org/10.14202/vetworld.2018.1-4>
- 6 • Saminathan M, Gopalakrishnan A, Latchumikanthan A, Milton AAP, Aravind M, Dhama K, Singh R (2015). Histopathological and parasitological study of blood-sucking *Haemonchus contortus* infection in sheep. Adv. Anim. Vet. Sci. 3(2): 99. <https://doi.org/10.14737/journal.aavs/2015/3.2.99.108>
- 34 • Somvanshi R, Rao JR (2009). Necropsy techniques and necropsy conference manual. Indian Vet. Res. Inst. Izatnagar-India. pp. 1-10.
- Soulsby E JL (1982). Helminths, arthropods and protozoa of domesticated animals. 7<sup>th</sup> ed. London. The English Book Society and Bailliere, Tindall.
- Stear MJ, Murray M (1994). Genetic resistance to parasitic disease: particularly of resistance in ruminants to gastrointestinal nematodes. Vet. Parasitol. 54(1-3): 161-176.
- 25 [https://doi.org/10.1016/0304-4017\(94\)90089-2](https://doi.org/10.1016/0304-4017(94)90089-2)
- Taraschewski H (2000). Host-parasite interactions in acanthocephala: a morphological approach. Adv. Parasitol. 46(1): 1-179. [https://doi.org/10.1016/S0065-308X\(00\)46008-2](https://doi.org/10.1016/S0065-308X(00)46008-2)
- 14 • Tariq KA, Chishti MZ, Ahmad F, Shawl AS (2009). Anthelmintic activity of extracts of *Artemisia absinthium* against ovine nematodes. Vet. Parasitol. 160(1-2): 83-88. <https://doi.org/10.1016/j.vetpar.2008.10.084>
- 4 • Urquhart GM, Armour JL, Duncan JL, Dunn AM, Jennings FW (1996). Veterinary parasitology longman scientific and technical. Paper. pp. 264.
- Vatta AF, Letty BA, Van der Linde MJ, Van Wijk EF, Hansen JW, Krecsek RC (2001). Testing for clinical anaemia caused by *Haemonchus* spp. in goats under resource-poor conditions in South Africa using an eye colour chart developed for sheep. Vet. Parasitol. 99(1): 1-14. [https://doi.org/10.1016/S0304-4017\(01\)00446-0](https://doi.org/10.1016/S0304-4017(01)00446-0)
- Widayati I, Nurhayati D (2013). Evaluation of fresh extracts and infusion of kebar grass (*Biophytum petersianum* Klotzsch) as anthelmintic against *Ascaridia galli* *in vitro*. <https://onsearch.id/Record/IOS1.INLIS.000000000623513>.



# In vivo Efficacy of Biophytum petersianum on Haemonchus contortus in Goats

## ORIGINALITY REPORT

**34%**

SIMILARITY INDEX

**29%**

INTERNET SOURCES

**29%**

PUBLICATIONS

**25%**

STUDENT PAPERS

## PRIMARY SOURCES

<b>1</b>	<b>Submitted to Mansoura University</b> Student Paper	<b>3%</b>
<b>2</b>	<b>Submitted to Associatie K.U.Leuven</b> Student Paper	<b>1%</b>
<b>3</b>	<b><a href="http://ejournal.undip.ac.id">ejournal.undip.ac.id</a></b> Internet Source	<b>1%</b>
<b>4</b>	<b>Venus Alma Appel Uruburu, Liseth Andrea Quiroz Acosta, Daniel Alejandro Noguera Ortega. "Aplicación del método Famacha© en dos tipos de explotación caprina en Popayán (Cauca, Colombia)", Revista de Medicina Veterinaria, 2017</b> Publication	<b>1%</b>
<b>5</b>	<b><a href="http://www.jherpmedsurg.com">www.jherpmedsurg.com</a></b> Internet Source	<b>1%</b>
<b>6</b>	<b>Submitted to Central Queensland University</b> Student Paper	<b>1%</b>
<b>7</b>	<b>Submitted to University of Zululand</b>	

1%

8

Iveta A. Kyriánová, Jaroslav Vadlejch, Oldřich Kopecký, Iva Langrová. "Seasonal dynamics of endoparasitic infections at an organic goat farm and the impact of detected infections on milk production", Parasitology Research, 2017

Publication

1%

9

[academicjournals.org](http://academicjournals.org)

Internet Source

1%

10

[www.myfoodresearch.com](http://www.myfoodresearch.com)

Internet Source

1%

11

Submitted to Chester College of Higher Education

Student Paper

1%

12

[www.scopemed.org](http://www.scopemed.org)

Internet Source

1%

13

Submitted to Charles Sturt University

Student Paper

1%

14

[ijgrr.org](http://ijgrr.org)

Internet Source

1%

15

[jurnal.ugm.ac.id](http://jurnal.ugm.ac.id)

Internet Source

1%

16

Antoine Rostang, Jacques Devos, Christophe Chartier. "Review of the Eprinomectin effective

1%



doses required for dairy goats: Where do we go from here?", Veterinary Parasitology, 2020

Publication

17

Abbas Imani-Baran, Jalal Abdollahi, Hamid Akbari, Saeed Jafarirad, Sajjad Moharramnejad. "Anthelmintic activity of crude powder and crude aqueous extract of *Trachyspermum ammi* on gastrointestinal nematodes in donkey (*Equus asinus*): An in vivo study", Journal of Ethnopharmacology, 2020

Publication

1%

18

Submitted to Higher Education Commission Pakistan

Student Paper

1%

19

e-sciencecentral.org

Internet Source

1%

20

Submitted to University of Aberdeen

Student Paper

1%

21

Saeed El-Ashram, Ibrahim Al Nasr, Fathi Abouhajer, Maged El-Kemary et al. "Microbial community and ovine host response varies with early and late stages of *Haemonchus contortus* infection", Veterinary Research Communications, 2017

Publication

1%

22

Abdelghany HEFNAWY, Mahmoud Atef Youssef

1%

HELAL, Ahmed SABEK, Saad SHOUSHA.  
"Clinical, behavioral and biochemical alterations  
due to shearing stress in Ossimi sheep", Journal  
of Veterinary Medical Science, 2018

Publication

23

Submitted to University of Queensland

Student Paper

1%

24

Submitted to Scotland's Rural College

Student Paper

1%

25

Jesús S. Hernández-Orts, Francisco E.  
Montero, Néstor A. García, Enrique A. Crespo  
et al. "Transmission of *Corynosoma australe*  
(Acanthocephala: Polymorphidae) from fishes to  
South American sea lions *Otaria flavescens* in  
Patagonia, Argentina", Parasitology Research,  
2019

Publication

1%

26

[www.jmbfs.org](http://www.jmbfs.org)

Internet Source

1%

27

[krishikosh.egranth.ac.in](http://krishikosh.egranth.ac.in)

Internet Source

1%

28

[ijarm.com](http://ijarm.com)

Internet Source

<1%

29

Pathak A.K.. "Potential of Using Condensed  
Tannins to Control Gastrointestinal Nematodes  
and Improve Small Ruminant Performance",

<1%



# International Journal of Molecular Veterinary Research, 2013

Publication

30

[www.ajol.info](http://www.ajol.info)

Internet Source

<1%

31

160.99.41.2

Internet Source

<1%

32

Javad Javanbakht, Ehsan Hosseini, Shadi Mousavi, Mehdi Aghamohammad Hassan et al. "Evaluation of two Iranian domestic ovine breeds for their pathological findings to gastrointestinal infection of *Haemonchus contortus*", Journal of Parasitic Diseases, 2013

Publication

<1%

33

[prp.hec.gov.pk](http://prp.hec.gov.pk)

Internet Source

<1%

34

Dharmendra Kanojiya, Daya Shanker, Vikrant Sudan, Amit Kumar Jaiswal, Rahul Parashar. " Assessment of and anthelmintic potential of extracts of bulb against naturally occurring ovine gastrointestinal nematodiosis ", Veterinary Quarterly, 2015

Publication

<1%

35

Purwaningsih Purwaningsih, Noviyanti Noviyanti, Priyo Sambodo. "INFESTASI CACING SALURAN PENCERNAAN PADA

<1%

KAMBING KACANG PERANAKAN ETTAWA DI  
KELURAHAN AMBAN KECAMATAN  
MANOKWARI BARAT KABUPATEN  
MANOKWARI PROVINSI PAPUA BARAT",  
JURNAL ILMIAH PETERNAKAN TERPADU,  
2017

Publication

36

[etd.lsu.edu](http://etd.lsu.edu)

Internet Source

<1%

37

[espace.curtin.edu.au](http://espace.curtin.edu.au)

Internet Source

<1%

38

Richard Komakech, Youngmin Kang.  
"Ethnopharmacological potential of African  
cherry [*Prunus africana*]", *Journal of Herbal  
Medicine*, 2019

Publication

<1%

39

[pt.scribd.com](http://pt.scribd.com)

Internet Source

<1%

40

Ibrahim Aly, Essam H. Ibrahim, Rabab S.  
Hamad, Hoda E. L. Sayed et al. "The Protective  
Role of Toll-Like Receptor Agonist  
Monophosphoryl Lipid A Against Vaccinated  
Murine Schistosomiasis", *Acta Parasitologica*,  
2020

Publication

<1%

41

[espace.library.uq.edu.au](http://espace.library.uq.edu.au)

Internet Source

<1%



- 
- 42 [doaj.org](http://doaj.org)  
Internet Source <1%
- 
- 43 Chinnaperumal Kamaraj. "Ovicidal and larvicidal activity of crude extracts of Melia azedarach against Haemonchus contortus (Strongylida)", Parasitology Research, 02/23/2010  
Publication <1%
- 
- 44 [recentscientific.com](http://recentscientific.com)  
Internet Source <1%
- 
- 45 E. BARRAU. "Effect of bioactive compounds from Sainfoin ( Onobrychis viciifolia Scop.) on the in vitro larval migration of Haemonchus contortus : role of tannins and flavonol glycosides", Parasitology, 06/16/2005  
Publication <1%
- 
- 46 [www.mdpi.com](http://www.mdpi.com)  
Internet Source <1%
- 
- 47 [ejournalfpikunipa.ac.id](http://ejournalfpikunipa.ac.id)  
Internet Source <1%
- 
- 48 [www.macvetrev.mk](http://www.macvetrev.mk)  
Internet Source <1%
- 
- 49 P. A. Pacheco, L. N. C. Rodrigues, J. F. S. Ferreira, A. C. P. Gomes et al. "Inclusion complex and nanoclusters of cyclodextrin to increase the solubility and efficacy of <1%

# albendazole", Parasitology Research, 2018

Publication

50

[www.dovepress.com](http://www.dovepress.com)

Internet Source

<1%

51

Tadesse Eguale. "Efficacy of albendazole against nematode parasites isolated from a goat farm in Ethiopia: relationship between dose and efficacy in goats", Tropical Animal Health and Production, 02/05/2009

Publication

<1%

52

Esmaeilnejad, B.. "Evaluation of antioxidant status and oxidative stress in sheep naturally infected with Babesia ovis", Veterinary Parasitology, 20120430

Publication

<1%

53

[search.crossref.org](http://search.crossref.org)

Internet Source

<1%

54

[livrepositary.liverpool.ac.uk](http://livrepositary.liverpool.ac.uk)

Internet Source

<1%

55

146.230.128.141

Internet Source

<1%

56

B. Santoso, A. Kilmaskossu, P. Sambodo. "Effects of saponin from Biophytum petersianum Klotzsch on ruminal fermentation, microbial protein synthesis and nitrogen utilization in goats", Animal Feed Science and Technology,

<1%

2007

Publication

---

57

[b-ok.org](http://b-ok.org)

Internet Source

<1%

---

58

[epdf.pub](http://epdf.pub)

Internet Source

<1%

---

59

Submitted to Harper Adams University College

Student Paper

<1%

---

60

H. M. Burrow. "Importance of adaptation and genotype × environment interactions in tropical beef breeding systems", animal, 2012

Publication

<1%

---

61

[mro.massey.ac.nz](http://mro.massey.ac.nz)

Internet Source

<1%

---

62

Cheryl Lans. "Validation of ethnoveterinary medicinal treatments", Veterinary Parasitology, 2011

Publication

<1%

---

63

E. Rakhshandehroo, M. Asadpour, S. H. Malekpour, A. Jafari. " The anthelmintic effects of five plant extracts on the viability of larvae ", Equine Veterinary Education, 2017

Publication

<1%

---

64

[www.pjbs.org](http://www.pjbs.org)

Internet Source

<1%

---

65

M. Worku, R. Franco, J.H. Miller. "Evaluation of the Activity of Plant Extracts in Boer Goats", American Journal of Animal and Veterinary Sciences, 2009

Publication

---

&lt;1%

66

Eguale, T.. "In vitro and in vivo anthelmintic activity of crude extracts of Coriandrum sativum against Haemonchus contortus", Journal of Ethnopharmacology, 20070404

Publication

---

&lt;1%

67

hdl.handle.net

Internet Source

---

&lt;1%

68

H. Hoste, J.F.J. Torres-Acosta, J. Quijada, I. Chan-Perez, M.M. Dakheel, D.S. Kommuru, I. Mueller-Harvey, T.H. Terrill. "Interactions Between Nutrition and Infections With Haemonchus contortus and Related Gastrointestinal Nematodes in Small Ruminants", Elsevier BV, 2016

Publication

---

&lt;1%

69

Fernando Horcajada-Sánchez, Álvaro Navarro-Castilla, Mariana Boadella, Isabel Barja. "Influence of livestock, habitat type, and density of roe deer (*Capreolus capreolus*) on parasitic larvae abundance and infection seroprevalence in wild populations of roe deer from central Iberian Peninsula", Mammal Research, 2018

&lt;1%



70

José Manuel Molina, Yeray Isidoro Hernández, Sergio Martín, Otilia Ferrer, Francisco Rodríguez, Antonio Ruiz. " Immune response in goats vaccinated with thiol-binding proteins from ", *Parasite Immunology*, 2018

Publication

<1%

---

71

Aida C. Cala, Jorge F. S. Ferreira, Ana Carolina S. Chagas, Javier M. Gonzalez et al. "Anthelmintic activity of *Artemisia annua* L. extracts in vitro and the effect of an aqueous extract and artemisinin in sheep naturally infected with gastrointestinal nematodes", *Parasitology Research*, 2014

Publication

<1%

---

72

Submitted to iGroup

Student Paper

<1%

---

73

K.A. Tariq, M.Z. Chishti, F. Ahmad, A.S. Shawl, M.A. Tantray. " Evaluation of anthelmintic activity of against gastrointestinal nematodes of sheep ", *Journal of Helminthology*, 2008

Publication

<1%

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

# In vivo Efficacy of *Biophytum petersianum* on *Haemonchus contortus* in Goats

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---