

# Response of Broiler Chickens to Triticale-Based Diets Supplemented with Microbial Enzymes (2. Microbial Profiles and Activities)

*by A Widodo*

---

**Submission date:** 24-Jun-2020 03:03AM (UTC-0700)

**Submission ID:** 1348991986

**File name:** Article\_Edi-5.pdf (398.96K)

**Word count:** 6405

**Character count:** 31855



## Response of Broiler Chickens to Triticale-Based Diets Supplemented with Microbial Enzymes (2. Microbial Profiles and Activities)

Widodo AE<sup>1,2</sup>, Nolan JV<sup>1</sup>, Akter M<sup>1,3</sup>, O'Neill HM<sup>4</sup> & Iji PA<sup>1</sup>

<sup>1</sup>School of Environmental and Rural Science, University of New England, Armidale, Australia

<sup>2</sup>Department of Animal Science, University of Papua, Manokwari - West Papua, Indonesia

<sup>3</sup>Department of Dairy and Poultry Science, Chittagong Veterinary and Animal Sciences University, Chittagong, Bangladesh

<sup>4</sup>AB Vista, 3 Woodstock Court, Blenheim Road, Marlborough Business Park, Marlborough, UK

Poultry Science Journal 2018, 6(1): 41-50

### Keywords

Triticale  
Phytase  
Xylanase  
Broiler chicken  
Microbial profiles

### Corresponding author

Paul Ade Iji  
[pauladeiji@gmail.com](mailto:pauladeiji@gmail.com)

### Article history

Received: August 17, 2017  
Revised: December 20, 2017  
Accepted: January 24, 2018

### Abstract

The microbial profiles and activities of microbes in the digesta from chickens on diets containing Bogong or Canobolas with or without inclusion of microbial enzymes were investigated on the 384 day-old male Ross 308 broiler chickens. There was a reduction in population of *C. perfringens* ( $P < 0.01$ ) of the ileum as a result of interaction between cultivar and xylanase. On the other hand, in the caeca, the population of entero bacteria were reduced ( $P < 0.01$ ) by the interaction between grain and phytase, as well as the interaction between grain, xylanase and phytase. Also, the inclusion of xylanase reduced the population of *C. perfringens* in the caeca. The cecal concentration of acetic acid ( $P < 0.05$ ) and butyric plus isobutyric acid ( $P < 0.01$ ) was increased in birds received Bogong and Canobolas diets. Supplementation of xylanase in Bogong diets increased (grain x xylanase,  $P < 0.05$ ) the acetic acid and lactic acid concentration in caeca while the reversed was the case for Canobolas diets. The ileal and cecal pH was not affected by grain, the inclusion of xylanase and phytase or interactions between these factors. It can be concluded that diets containing Bogong or Canobolas diets supplemented with phytase and xylanase influenced the microbial profile and their activities in gastrointestinal tract which may be due to the variation in nutrient content of these two cultivars.

### Introduction

In the trials conducted by South Australian Research and Development Institute, Bogong and Canobolas were found to be the highest yielding new cultivars of triticale (Crouch and Saunders, 2009). In the previous work (Widodo *et al.*, 2015), it shown that diets based on these two new cultivars were supported excellent production of broiler chickens. The primary constraint to the use of triticale in poultry diets is the presence of non-starch polysaccharides (NSP), especially xylans and arabinoxylans, as well as phytate, which together reduce nutrient digestibility (Antoniou and Marquardt, 1981).

The inclusion of xylanase and phytase in diets based on Bogong and Canobolas has already been shown to improve the nutritive value of such diets and improve the productivity of broiler chickens (Widodo *et al.*, 2011). Feed enzymes are known to reduce the bacterial activity in the ileum by reducing the amount of nutrient available for microbial fermentation (Silva and Smithard, 2002). Jamroz *et al.* (2002) suggested that the end-products of microbial fermentation (e.g., short-chain fatty acids; SCFA) in the chicken intestine may contribute energy to the host bird. In addition, Ricke (2003) and van der Wielen *et al.* (2000) reported that dietary

enzymes may play an important role in regulating the gastrointestinal tract (GIT) microbial population. The concentrations of SCFA and lactic acid in the GIT depend on the cereal type and feed enzymes used (Jamroz *et al.*, 2002; Silva and Smithard, 2002; Józefiak *et al.*, 2004a; 2004b) and reflect to a certain extent the activity of the resident microflora (Engberg *et al.*, 2002).

The aim of this work was to investigate how feeding of diets based on triticale, cultivar

Bogong and Canobolas, and supplemented with xylanase and phytase influence the gastrointestinal ecosystems of broiler chickens in terms of microbial profiles, concentrations of organic acids and pH.

## Materials and Methods

### Experimental design and bird management

The experiment was approved by the Animal Ethics Committee of the University of New England, Australia (Approval No. AEC 10/098).

**Table 1.** Ingredients and nutrient composition (g/kg) of dietary treatments

Ingredients	B	BX	BP	BXP	C	CX	CP	CXP
Bogong	650.0	650.0	650.0	650.0	-	-	-	-
Canobolas	-	-	-	-	650.0	650.0	650.0	650.0
Soybean Meal	190.0	190.0	190.0	190.0	190.0	190.0	190.0	190.0
Soycomil K	69.4	69.4	69.4	69.4	61.3	61.2	61.2	61.2
L-Threonine	1.8	1.8	1.8	1.8	1.9	1.9	1.9	1.9
L-Lysine HCl	4.8	4.8	4.8	4.8	5.3	5.3	5.3	5.3
DL-Methionine	2.6	2.6	2.6	2.6	3.0	3.0	3.0	3.0
Sunflower oil	35.7	35.7	35.5	35.3	42.6	42.6	42.5	42.4
Limestone	18.1	18.1	18.1	18.1	18.1	18.1	18.1	18.1
Dical. P	13.8	13.8	13.8	13.8	13.8	13.8	13.8	13.8
Common Salt	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Choline Cl-70%	1.6	1.6	1.6	1.6	1.5	1.5	1.5	1.5
Xylanase	-	0.1	-	0.1	-	0.1	-	0.1
Phytase	-	-	0.2	0.2	-	-	0.2	0.2
Premix <sup>1</sup>	2.6	2.6	2.6	2.6	2.5	2.5	2.5	2.5
TiO <sub>2</sub>	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
<i>Nutrient composition</i>								
ME (Kcal/kg)	3033	3033	3033	3033	3081	3081	3081	3081
Crude protein	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0
Crude fat	53.8	53.7	53.6	53.5	59.4	59.3	59.2	59.1
Crude fiber	25.5	25.5	25.5	25.5	25.2	25.2	25.2	25.2
Lysine	15.0	15.0	15.0	15.0	15.0	15.0	15.0	15.0
Methionine	6.1	6.1	6.1	6.1	6.5	6.5	6.5	6.5
Met + Cys	10.5	10.8	10.8	10.8	11.3	11.3	11.3	11.3
Calcium	11.1	11.1	11.1	11.1	11.1	11.1	11.1	11.1
Available P	5.2	5.2	5.2	5.2	5.6	5.6	5.6	5.6
Sodium	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
Chlorine	5.6	5.6	5.6	5.6	3.4	3.4	3.4	3.4

<sup>1</sup> Supplied per kg of diet (mg): vitamin A (as all-trans retinol): 3.6; cholecalciferol: 0.09; vitamin E (as d- $\alpha$ -tocopherol): 44.7; vitamin K<sub>3</sub>: 2.0; thiamine: 2.0; riboflavin: 6.0; pyridoxine hydrochloride: 5.0; vitamin B<sub>12</sub>: 0.2; biotin: 0.1; niacin: 50.1; D-calcium pantothenate: 12.0; folic acid: 2.0; Mn: 80.0; Fe: 60.0; Cu: 8.0; I: 1.0; Co: 0.3; and Mo: 1.0.

### Bird management and treatments

The experiment was conducted in a 2 × 2 × 2 factorial arrangement. Eight experimental diets were formulated to study the effect of 2 cultivars of high-yielding triticale (Bogong and Canobolas), with or without xylanase and with or without phytase on intestinal microbial profile and activities. The experimental diets contained 65% of either Bogong or Canobolas. The microbial phytase, Quantum<sup>®</sup> 2500, which is a 6-phytase from *E. coli* (AB Vista, Marlborough,

UK) was added to supply 500 FTU per kg diet. The xylanase preparation, Econase<sup>®</sup> XT, which contains thermostable endo-1,4-beta-xylanase, produced by *Trichoderma reesei*, was added to supply 160,000 BXU of xylanase activity. In all diets, titanium dioxide (TiO<sub>2</sub>) was incorporated as an indigestible marker to enable assessment of nutrient digestibility. All diets were pelleted after mixing and fed up to 21 days of age. The experimental diets were formulated to meet the minimum Aviagen breeder recommendations

(Aviagen, 2007). The details of the ingredients and nutrients composition of all diets are described in Table 1. The dietary treatments were as follows: a diet based on Bogong without any enzymes (B); Bogong with the inclusion of xylanase (BX); Bogong with the inclusion of phytase (BP); Bogong with the inclusion of xylanase and phytase (BXP); Canobolas without enzymes (C); Canobolas with the inclusion of xylanase (CX); Canobolas with the inclusion of phytase (CP), and Canobolas with the inclusion of xylanase and phytase (CXP).

A total of 384 day-old male Ross 308 broiler chicks (Baiada Poultry Pty. Ltd, Tamworth, NSW, Australia), weighing  $41.30 \pm 0.35$  g, were randomly assigned to eight treatments. Each of the treatments had six replicates, with 8 birds per replicate cage. The experimental chickens were raised in battery brooders,  $60 \times 42 \times 23$  cm, set up in a climate-controlled room. The birds were initially brooded at a temperature of  $34^\circ\text{C}$ , which was reduced gradually to  $24 \pm 1^\circ\text{C}$  at 21 days of age when the feeding trial ended. Light was provided for 18 h per day throughout the trial. The birds had free access to water and feed.

On day 21, three birds from each cage, were randomly selected, weighed and killed by cervical dislocation. The abdominal cavity was opened and the small intestine was ligated and removed. The contents of the ileum and caeca were squeezed and collected in plastic containers and pooled by cage for the determination of SCFA. Around 1 g of ileal and cecal contents were also collected separately into prepared McCartney bottles containing anaerobic broth for the enumeration of microbial populations. The McCartney bottles and plastic containers containing digesta samples for microbial profiles and SCFA concentrations, respectively, were kept at  $-20^\circ\text{C}$  until they were analyzed.

#### Gut microbial population

Approximately 1 g of sample of either ileal or cecal digesta was placed into a pre-weighed McCartney bottle containing 8 mL of anaerobic broth and 1 mL of reducing (R) solution. After addition of sample to the anaerobic broth, the bottle with broth and sample was weighed again. Once the sample was weighed, the first dilution of the sample was calculated by subtracting the weight of the bottle and the broth plus the R solution from the final weight of the bottle containing the sample (this is the sample weight),

and dividing the sample weight by the broth plus R solution. This calculation yield is the proportion of the original sample per mL of solution in the bottle. The suspension was homogenized for 2 min in  $\text{CO}_2$ -flushed plastic bags using a bag mixer (Interscience, St. Norm, France) and then serially diluted in 10-fold increments in anaerobic broth according to the procedure described by Miller and Wolin (1974) and Engberg *et al.* (2004). A 1 mL aliquot of the homogenized suspension was then transferred into 9 mL of anaerobic broth and serially diluted from  $10^{-1}$  to  $10^{-5}$  for ileum samples and  $10^{-1}$  to  $10^{-6}$  for the cecal samples. From the last three diluted samples, 0.1 mL of each sample was plated on the appropriate medium (10 mL) for enumeration of microbial populations.

The enumeration of anaerobic bacteria was done by following the method described by Miller and Wolin (1974) and Engberg *et al.* (2004). Total anaerobic bacteria were grown in anaerobic roll tubes containing 3 mL of Wilkins-Chalgren anaerobic agar (Oxoid, CM 0619) incubated at  $39^\circ\text{C}$  for 7 days. Lactic acid bacteria were enumerated after incubation on MRS agar (Oxoid, CM0361) under anaerobic conditions at  $39^\circ\text{C}$  for 48 h. Lactobacilli were enumerated on Rogossa agar (Oxoid, CM 0627) after anaerobic incubation at  $39^\circ\text{C}$  for 48 h. Enterobacteria were incubated on MacConkey agar (Oxoid, CM 0007) aerobically at  $39^\circ\text{C}$  for 24 h prior to counting. *Clostridium perfringens* (Cp) colonies were counted on Tryptose-Sulfite-Cycloserine and Shahidi-Ferguson Perfringens agar base (TSC & SFP) (Oxoid, CM 0587 OPSP) mixed with egg yolk emulsion (Oxoid, SR0047) and Perfringens (TSC) selective supplement (Oxoid, SR0088E) according to the pour-plate technique, where plates were overlaid with the same agar after spreading the inoculum and incubated anaerobically at  $39^\circ\text{C}$  for 24 h. An anaerobic Anaero Gen™ sachet (AN0025A, Oxoid Ltd, Hampshire, UK) was used to generate the anaerobic environment ( $<1\% \text{O}_2$  and  $9\text{--}13\% \text{CO}_2$ ) for all anaerobically incubated agar plates. After incubation, colonies formed on the respective media were carefully counted, converted into logarithmic equivalents ( $\log_{10}$ ) and expressed as numbers of colony-forming units (CFU) per gram of wet ileal or cecal digesta.

#### Measurement of organic acid

A modification of the analytical method described by Jensen *et al.* (1995) for the analysis of organic acids (SCFAs, lactic acid and succinic acid) concentrations was used. In general, frozen

ileal and cecal samples were thawed and homogenized by vigorous shaking. About 1 to 2 g of ileal and cecal digesta (wet weight) were accurately weighed into centrifuge tubes (placed on ice) and 1 mL internal standard (0.01 methyl butyric acid) was added and thoroughly mixed with a vortex mixer, followed by centrifugation at  $25,700 \times g$  at  $5^{\circ}\text{C}$  for 20 min in a Beckman model J2-21M Induction Drive Centrifuge with a JA-21 rotor. The supernatant (approximately 1 mL) was transferred into 8-mL vials (on ice). At this stage, the standards and blanks were prepared. One mL of the standard acid mixture was accurately transferred to the 8-mL vials. The blank was treated similarly (1 mL of Milli-Q water), and 0.1 mL of 0.1 M ethyl butyric acid was added to the standards and blank vials. Two mL of ether and 0.5 mL of concentrated HCl (36%) were then added to all the samples, the standard and the blank and the lids were tightened prior to vortexing for about 1 minute. After repeat vortexing, all the tubes were centrifuged at  $2060 \times g$  for 15 min at  $5^{\circ}\text{C}$ . Accurately, 400  $\mu\text{L}$  of the supernatant were transferred into gas chromatograph (GC) vials (2 mL) and mixed with 40  $\mu\text{L}$  of N-tert-butyltrimethylsilyl-N-methyl trifluoroacetamide (MTBSTFA) or 50  $\mu\text{L}$  MTBSTFA for the standard. The GC vials were tightly capped, vortexed and kept on a heating block at  $80^{\circ}\text{C}$  for 20 min and then left at room temperature for 48 h. After 48 h, concentrations of the different organic acids were determined, using a Varian CP3400 CX gas chromatograph (Varian Analytical Instruments, Palo Alto, CA, USA).

#### Intestinal pH

The pH of ileal and cecal digesta was measured on fresh samples collected at 21 d of age. The pH was determined by the modified procedure of Corrier *et al.* (1990). Around 1 g of the content was diluted in 9 mL of cold distilled MilliQ water. The suspension was mixed thoroughly with a stirrer and the pH was determined by insertion of a glass electrode (EcoScan 5/6 pH meter, Eutech Instruments Pte Ltd., Singapore).

#### Statistical analysis

All data were analyzed by ANOVA using the general linear model (GLM) procedure of Minitab® Version 16 (Minitab, 2010) for the main

factors (cultivar, xylanase, and phytase) and the interactions between these three factors. The significance of difference between means was determined by Fisher's least significant difference (LSD) test, for which the significance level was set at  $P < 0.05$ .

## Results

### Gut microflora

The populations of total anaerobic bacteria (TAB), lactobacilli, and Enterobacteria in the ileal digesta were not affected ( $P > 0.05$ ) by grain cultivar, inclusion of xylanase or phytase (Table 2). An interaction between grain cultivar, xylanase and phytase was significant ( $P < 0.01$ ) in the case of the population of lactic acid bacteria (LAB), tending to rise with the inclusion of both enzymes. A grain  $\times$  xylanase interaction ( $P < 0.01$ ) was noticed on *C. perfringens* population, which in Canobolas-based diet was higher than in Bogong-based diet. Furthermore, there was an interaction ( $P = 0.067$ ) between xylanase and phytase on the population of *C. perfringens*, although there were opposite trends in Bogong- and Canobolas-based diets. In Bogong-based diets, the population of *C. perfringens* was less than on the control diet while it was higher in Canobolas-based diet.

In the cecal digesta, the grain, xylanase and the phytase inclusion did not affect the population of TAB, LAB, and lactobacilli (Table 3). Meanwhile, the inclusion of xylanase decreased ( $P < 0.01$ ) the population of *C. perfringens*. An interaction between grain and phytase was noticed in the population of enterobacteria ( $P < 0.05$ ) and *C. perfringens* ( $P = 0.055$ ). The population of enterobacteria was less in Bogong diet plus phytase than the Bogong diet without enzyme, however; it was opposite in the Canobolas diet, which the Canobolas diet plus phytase had higher population of enterobacteria than the Canobolas diet without enzyme. An interaction ( $P < 0.01$ ) between grain, xylanase and phytase inclusion indicated that the population of enterobacteria that in Bogong diet decreased the population, on the contrary, it increased in Canobolas diet. The interaction between grain, xylanase and phytase inclusion tended ( $P = 0.074$ ) to decrease the population of *C. perfringens* in both diets with the inclusion of xylanase and phytase.

**Table 2.** Bacterial counts (log<sub>10</sub> CFU/g digesta) in ileal digesta of broiler chickens triticale-based diets with or without enzyme supplementation<sup>1</sup>

Treatments			Total Anaerobic	Lactic acid	Lactobacilli	Enterobacteria	C. <i>perfringens</i>
Grain	Xyl <sup>2</sup>	Phy <sup>3</sup>					
Bogong	-	-	7.1	8.0 <sup>abc</sup>	7.1	4.6	4.3 <sup>abc</sup>
Bogong	+	-	7.1	8.2 <sup>ab</sup>	7.3	4.7	4.3 <sup>abc</sup>
Bogong	-	+	7.0	8.1 <sup>abc</sup>	7.1	4.8	4.5 <sup>ab</sup>
Bogong	+	+	7.1	7.8 <sup>bc</sup>	7.1	4.8	3.9 <sup>c</sup>
Canobolas	-	-	7.1	8.5 <sup>a</sup>	7.1	4.6	4.1 <sup>bc</sup>
Canobolas	+	-	7.1	7.6 <sup>c</sup>	7.1	4.8	4.8 <sup>a</sup>
Canobolas	-	+	7.0	8.2 <sup>ab</sup>	7.1	4.8	4.2 <sup>bc</sup>
Canobolas	+	+	7.0	8.4 <sup>a</sup>	7.1	5.0	4.5 <sup>ab</sup>
Pooled SEM <sup>4</sup>			0.08	0.07	0.08	0.04	0.07
Source of variation			Significance of treatment effect				
Grain			ns	ns	ns	ns	ns
Xylanase			ns	ns	ns	ns	ns
Phytase			ns	ns	ns	ns	ns
Grain × Xylanase			ns	ns	ns	ns	**
Grain × Phytase			ns	ns	ns	ns	ns
Xylanase × Phytase			ns	ns	ns	ns	0.067
Grain × Xylanase × Phytase			ns	**	1	ns	ns

<sup>1</sup> Each value represents the mean of 6 replicates; <sup>2</sup>Xylanase; <sup>3</sup>Phytase; <sup>4</sup>SEM = Standard error of means.

<sup>a-c</sup> Values with unlike superscripts within each column are significantly different at \*\* $P < 0.01$ ; ns = not significant.

**Table 3.** Bacterial counts (log<sub>10</sub> CFU/g digesta) in caecal digesta of broiler chickens fed triticale-based diets with or without enzyme supplementation<sup>1</sup>

Treatments			Total Anaerobic	Lactic acid	Lactobacilli	Enterobacteria	C. <i>perfringens</i>
Grain	Xyl <sup>2</sup>	Phy <sup>3</sup>					
Bogong	-	-	8.8	9.4	7.8	8.1 <sup>a</sup>	6.1 <sup>a</sup>
Bogong	+	-	8.8	9.3	7.7	8.0 <sup>abc</sup>	5.8 <sup>bc</sup>
Bogong	-	+	8.7	9.4	7.9	7.8 <sup>bcd</sup>	6.0 <sup>ab</sup>
Bogong	+	+	8.7	9.3	7.9	7.8 <sup>abcd</sup>	5.9 <sup>abc</sup>
Canobolas	-	-	8.7	9.4	7.7	7.7 <sup>d</sup>	6.1 <sup>a</sup>
Canobolas	+	-	8.7	9.4	7.9	8.1 <sup>a</sup>	6.0 <sup>a</sup>
Canobolas	-	+	8.7	9.3	7.8	8.0 <sup>ab</sup>	6.0 <sup>a</sup>
Canobolas	+	+	8.7	9.3	7.8	7.8 <sup>cd</sup>	5.7 <sup>c</sup>
Pooled SEM <sup>4</sup>			0.05	0.04	0.04	0.04	0.03
Source of variation			Significance of treatment effect				
Grain			ns	ns	ns	ns	ns
Xylanase			ns	ns	ns	ns	**
Phytase			ns	ns	ns	ns	ns
Grain × Xylanase			ns	ns	ns	ns	ns
Grain × Phytase			ns	ns	ns	*	0.055
Xylanase × Phytase			ns	ns	ns	0.066	ns
Grain × Xylanase × Phytase			ns	ns	1	**	0.074

<sup>1</sup> Each value represents the mean of 6 replicates; <sup>2</sup>Xylanase; <sup>3</sup>Phytase; <sup>4</sup>SEM = Standard error of means.

<sup>a-c</sup> Values with unlike superscripts within each column are significantly different at \* $P < 0.05$ ; \*\* $P < 0.01$ ; ns = not significant.

### Short-chain fatty acids, lactic acid and succinic acid

The concentrations of formic, acetic, butyric, lactic, succinic acids and lactic acid of ileal are shown in Table 4. There was no significant difference in concentrations of formic, lactic and

succinic acids, but acetic acid concentration decreased ( $P < 0.01$ ) in birds on diets supplemented with phytase and was numerically reduced ( $P = 0.066$ ) by the inclusion of xylanase in both Bogong- and Canobolas-based diets.

**Table 4.** Concentrations of various short-chain acids ( $\mu\text{mol/g}$  wet digesta) in ileal contents of broiler chickens on triticale-based diets with or without enzyme supplementation<sup>1</sup>

Treatments						
Grain	Xyl <sup>2</sup>	Phy <sup>3</sup>	Formic	Acetic	Lactic	Succinic
Bogong	-	-	0.31	1.6 <sup>ab</sup>	22.6	0.35
Bogong	+	-	0.34	1.4 <sup>abc</sup>	17.1	0.24
Bogong	-	+	0.31	1.2 <sup>bc</sup>	16.0	0.83
Bogong	+	+	0.31	1.0 <sup>c</sup>	19.2	0.44
Canobolas	-	-	0.43	1.8 <sup>a</sup>	15.3	0.19
Canobolas	+	-	0.30	1.4 <sup>abc</sup>	11.9	0.34
Canobolas	-	+	0.37	1.2 <sup>bc</sup>	12.4	0.35
Canobolas	+	+	0.25	0.9 <sup>c</sup>	13.8	0.11
Pooled SEM <sup>4</sup>			0.015	0.08	1.08	0.07
Source of variation			Significance of treatment effect			
Grain			ns	ns	ns	ns
Xylanase			ns	0.066	ns	ns
Phytase			ns	**	ns	ns
Grain × Xylanase			ns	ns	ns	ns
Grain × Phytase			ns	ns	ns	ns
Xylanase × Phytase			ns	ns	ns	ns
Grain × Xylanase × Phytase			ns	ns	ns	ns

<sup>1</sup>Each value represents the mean of 6 replicates; <sup>2</sup>Xylanase; <sup>3</sup>Phytase; <sup>4</sup>SEM = Standard error of means.

\* < Values with unlike superscripts within each column are significantly different at \*\*  $P < 0.01$ ; ns = not significant.

**Table 5.** Concentration of various organic and mineral acids ( $\mu\text{mol/g}$  wet digesta) in caecal contents of broiler chickens on high-yielding triticale-based diet with and without enzymes<sup>1</sup>

Treatments								
Grain	Xyl <sup>2</sup>	Phy <sup>3</sup>	Acetic	Propionic	Butyric + Isobutyric	Valeric + Isovaleric	Lactic	Succinic
Bogong	-	-	76.2 <sup>c</sup>	7.2 <sup>bc</sup>	19.8 <sup>bc</sup>	1.6 <sup>b</sup>	9.1 <sup>ab</sup>	25.2
Bogong	+	-	133.3 <sup>ab</sup>	6.4 <sup>bc</sup>	30.8 <sup>ab</sup>	1.8 <sup>ab</sup>	19.4 <sup>a</sup>	38.5
Bogong	-	+	104.2 <sup>bc</sup>	9.8 <sup>ab</sup>	21.7 <sup>bc</sup>	2.0 <sup>ab</sup>	4.7 <sup>b</sup>	34.8
Bogong	+	+	169.3 <sup>a</sup>	12.5 <sup>a</sup>	37.2 <sup>a</sup>	2.7 <sup>a</sup>	18.7 <sup>a</sup>	51.2
Canobolas	-	-	103.8 <sup>bc</sup>	7.4 <sup>bc</sup>	13.4 <sup>c</sup>	1.6 <sup>b</sup>	7.4 <sup>ab</sup>	52.5
Canobolas	+	-	107.5 <sup>bc</sup>	4.6 <sup>c</sup>	18.6 <sup>c</sup>	1.1 <sup>b</sup>	1.2 <sup>b</sup>	30.5
Canobolas	-	+	100.6 <sup>bc</sup>	6.9 <sup>bc</sup>	17.9 <sup>c</sup>	1.6 <sup>b</sup>	2.0 <sup>b</sup>	33.7
Canobolas	+	+	98.2 <sup>bc</sup>	3.7 <sup>c</sup>	20.2 <sup>bc</sup>	1.2 <sup>b</sup>	1.2 <sup>b</sup>	35.7
Pooled SEM <sup>4</sup>			7.09	0.63	1.74	0.13	1.72	3.54
Source of variation			Significance of treatment effect					
Grain			ns	**	**	*	**	ns
Xylanase			*	ns	**	ns	ns	ns
Phytase			ns	ns	ns	ns	ns	ns
Grain × Xylanase			*	0.081	ns	0.058	*	ns
Grain × Phytase			ns	*	ns	ns	ns	ns
Xylanase × Phytase			ns	ns	ns	ns	ns	ns
Grain × Xylanase × Phytase			ns	ns	ns	ns	ns	ns

<sup>1</sup>Each value represents the mean of 6 replicates; <sup>2</sup>Xylanase; <sup>3</sup>Phytase; <sup>4</sup>SEM = Standard error of means.

\* < Values with unlike superscripts within each column are significantly different at \*  $P < 0.05$ ; \*\*  $P < 0.01$ ; ns = not significant.

#### Various organic and mineral acids concentration

In the caecal contents, the organic acids recovered were acetic, propionic, butyric and isobutyric, valeric and isovaleric, as well as lactic and succinic acids (Table 5). The grain factor affected ( $P < 0.01$ ) the concentration of propionic, butyric plus isobutyric and lactic acid, which the concentration of that acids was higher in Bogong


diets than in Canobolas diets. In addition, the concentration of valeric plus isovaleric acid was higher ( $P < 0.05$ ) in Bogong diets than in Canobolas diets. The inclusion of xylanase in both diets significantly increased acetic acid concentration ( $P < 0.05$ ) and butyric plus isobutyric concentration ( $P < 0.01$ ). There was no effect ( $P > 0.05$ ) of phytase inclusion on the concentration of organic acids.

A grain  $\times$  xylanase interaction ( $P < 0.05$ ) indicated that xylanase inclusion increased acetic acid and lactic acid concentration in Bogong diets but decreased in Canobolas diets. Meanwhile, the interaction between grain and xylanase ( $P = 0.081$ ) and ( $P = 0.58$ ) affected the concentration of propionic acid and valeric plus isovaleric acid, respectively. In addition, an interaction ( $P < 0.05$ ) between grain and phytase inclusion was noticed in the concentration of propionic acid, which in Bogong diet with phytase was higher than Bogong diet without enzyme, on the other hand, in Canobolas diet

plus phytase, the concentration was less than the Canobolas diet without enzyme.

#### Intestinal pH

There were no significant effects of grain, the inclusion of xylanase and phytase or interactions between these factors for both ileal and caecal pH (Fig. 1); however, the pH of the ceceal content tended ( $P = 0.07$ ) to be higher on Canobolas than Bogong-based diets. Furthermore, the caecal pH was numerically reduced by the inclusion of xylanase as well as the combination of supplemental xylanase and phytase.



**Figure 1.** pH of ileal and caecal contents of broiler chickens on triticale-based diets with or without enzymes supplementation.

Bogong without any enzymes (B); Bogong with the inclusion of xylanase (BX); Bogong with the inclusion of phytase (BP); Bogong with the inclusion of xylanase and phytase (BXP); Canobolas without enzymes (C); Canobolas with the inclusion of xylanase (CX); Canobolas with the inclusion of phytase (CP) and Canobolas with the inclusion of xylanase and phytase (CXP)

#### Discussion

##### Microbial profiles and activities

In this study, the interaction between grain, xylanase and phytase significantly influenced the count of LAB in ileum. Birds fed the Canobolas diets supplemented with both enzymes showed the highest count of LAB population in the ileum compared to Bogong diet with both enzymes. This result indicates that grain cultivar rather than enzyme type influenced the LAB population in ileum and similar observation was reported in a previous study where NSP composition in grain ingredients influenced the GIT microbial profile (Apajalahti *et al.*, 2004). The presence of high concentrations of both soluble and insoluble NSP, especially arabinose and xylose in Canobolas variety may be the reason for this increase LAB concentration. Soluble NSP increases the digesta viscosity whereas insoluble NSP reduce digesta passage rate by slowing down intestinal motility and

hindering the access of endogenous enzymes for respective substrate (Bedford and Schulze, 1998) may lead to increase in microbial population in the GIT (Hubener *et al.*, 2002). In contrast, the lowest number of LAB population in the CX diets than Canobolas diet with or without both enzymes is partly consistent with results of other studies (Engberg *et al.*, 2004) where xylanase supplementation not only increased LAB population in ileum but also the lactic acid concentration in caeca. However, this is not completely true for the present study as the concentration of lactic acid in caeca was low in the all Canobolas diets than Bogong diets. The reason behind this is not clear.

In the caeca, the population of Enterobacter was reduced in birds fed Bogong diets with phytase supplementation than those fed the Canobolas diet with phytase and this can be explained by the variation in P level in



forementioned diets. It has been reported that phytase can regulate bacterial growth in intestine but its effect strictly depends on dietary Ca and P levels or Ca:P ratio (Ptak *et al.*, 2015). A study reported that supplementation of phytase to low P diet improved the absorption of available phosphorus and Ca in distal ileum and reduced the availability of Ca and P in the caeca (Metzler-Zebeli *et al.*, 2008). Because of the involvement of Ca and P in bacterial growth, the reduction of these two minerals consequently reduces the proliferation of certain bacteria (Metzler-Zebeli *et al.*, 2010).

Regardless of grain cultivar, phytase supplementation significantly reduced the acetic acid concentration in ileum. In contrast, Ptak *et al.* (2015) reported that phytase supplementation increased the concentration of lactic and acetic acids in ileum only with a diet containing reduced level of Ca (8 g/kg) and digestible P levels (2.69 g/kg). It has been reported that formation of calcium phosphate complex (Ptak *et al.*, 2015) or low P concentration could reduce the fermentation process in the GIT (Komisarczuk *et al.*, 1987; Smulikowska *et al.*, 2010). Therefore, it is possible that the highest level of dietary Ca (11.1 g/kg) and P (5.2 and 5.6 g/kg in Bogong and Canobolas, respectively) in the diets of the present study could facilitate the formation of Ca-phosphate complex, resulting in depletion of available P due to impaired phytase activity and subsequently reduce the microbial fermentation process in the ileum. The concentration of lactic acid in caeca was increased in birds fed Bogong diets with xylanase or xylanase + phytase than on Canobolas diets. The presence of high concentration of lactic acid in caeca indicates that xylanase supplementation did not completely digest the NSP, therefore, undigested carbohydrate from ileum would reach the caeca and be fermented by caecal microflora (Choct *et al.*, 1999; Bedford, 2000).

Although SCFA produced in the caeca can yield only relatively small amounts of energy, they can provide other benefits. It has been reported that high fermentation activity in the caeca of chickens correlates with a lower pH, which has the potential to inhibit some pathogenic bacteria (Russell, 1992; van der Wielen *et al.*, 2000). In addition, McHan and Shotts (1993) reported the toxic effect of SCFA

on some enterobacteria such as *Salmonella typhimurium*. Using *in vitro* technique, they showed that the presence of SCFA could reduce the numbers of these bacteria by up to 50–80% but there was no such trend observed in the present study. In addition, the presence of certain SCFA in the chicken gut, such as lactic and acetic acids, have been associated with a decrease in the survival and adherence of *Salmonella enterica*, *E. coli* and *C. perfringens* (Engberg *et al.*, 2002; Engberg *et al.*, 2004; Bjerrum *et al.*, 2005). Józefiak *et al.* (2004b) reported that lactic acid is the main by-product of carbohydrate fermentation, which is produced by lactic acid bacteria. Lactic acid has been found to be important to broiler gut health and human food safety because its presence can inhibit enterobacteria (Bjerrum *et al.*, 2005). In this study, it was found that the population of enterobacteria and *C. perfringens*, which are considered to be risk groups, is generally smaller than that of lactobacilli that are regarded as beneficial bacteria; however, the increased population of enterobacteria and *C. perfringens* was undoubtedly affected by the supplemental enzymes on the diets. There were no symptoms of necrotic enteritis in this experiment that could be connected to the population of *C. perfringens*. Moreover, the mortality on this experiment was negligible, less than 1% and occurring in the first 7 d with no indications of infectious diseases.

### Conclusion

The lactic acid bacteria and *C. perfringens* in the ileum and enterobacteria and *C. perfringens* in the caeca were significantly affected by the dietary treatments. Furthermore, concentrations of organic acids in the digesta were affected by grain type and enzyme supplementation. The population of gram negative bacterial in the digesta was significantly reduced by the inclusion of xylanase and the combination of xylanase and phytase; however, the values were higher on the Bogong-based diets than on the Canobolas-based diets. It is not known if this difference contributed in any way to the difference in performance.

### Acknowledgements

This work was supported by AB Vista and the University of New England, Armidale Australia.

## References

- Antoniou T & Marquardt RR. 1981. Influence of rye pentosans on the growth of chicks. *Poultry Science*, 60: 1898-1904. DOI: 10.3382/ps.0601898
- Apajalahti J, Kettunen A & Graham H. 2004. Characteristics of the gastrointestinal microbial communities, with special reference to the chicken. *World's Poultry Science Journal*, 60: 223-232. DOI: 10.1079/WPS200415
- Aviagen. 2007. ROSS 308 BROILER: Nutrition Specification. Newbridge, Midlothian EH28 8SZ, Scotland, UK.
- Bedford MR. 2000. Removal of antibiotic growth promoters from poultry diets: implications and strategies to minimise subsequent problems. *World's Poultry Science Journal*, 56: 347-365. DOI: 10.1079/WPS20000024
- Bedford MR, & Schulze H. 1998. Exogenous enzymes for pigs and poultry. *Nutrition Research Review*, 11: 91-114. DOI: 10.1079/NRR19980007
- Bierrum L, Pedersen K & Engberg RM. 2005. The influence of whole wheat feeding on salmonella infection and gut flora composition in broilers. *Avian Disease*, 49: 9-15. DOI: 10.1637/7223-061504R
- Choct M, Hughes RJ & Bedford MR. 1999. Effects of a xylanase on individual bird variation, starch digestion throughout the intestine, and ileal and caecal volatile fatty acid production in chickens fed wheat. *British Poultry Science*, 40: 419-422. DOI: 10.1080/00071669987548
- Corrier DE, Hinton AJr, Ziprin RL, Beier RC & DeLoach JR. 1990. Effect of dietary lactose on cecal pH, bacteriostatic volatile fatty acids, and *Salmonella typhimurium* colonization of broiler chicks. *Avian Disease*, 34: 617-625. DOI: 10.2307/1591254
- Crouch J & Saunders R. 2009. Triticale Overview: New varieties lead in triticale trials. [http://www.sardi.sa.gov.au/crops/new\\_variety\\_agronomy/variety\\_evaluation/crop\\_harvest\\_reports/triticale/triticale\\_overview](http://www.sardi.sa.gov.au/crops/new_variety_agronomy/variety_evaluation/crop_harvest_reports/triticale/triticale_overview)
- Engberg RM, Hedemann MS & Jensen BB. 2002. The influence of grinding and pelleting of feed on the microbial composition and activity in the digestive tract of broiler chickens. *British Poultry Science*, 43: 569-579. DOI: 10.1080/0007166022000004480
- Engberg RM, Hedemann MS, Steinfeldt S & Jensen BB. 2004. Influence of whole wheat and xylanase on broiler performance and microbial composition and activity in the digestive tract. *Poultry Science*, 83: 925-938. DOI: 10.1080/0007166022000004480
- Jamroz D, Jakobsen K, Bach Knudsen KE, Wilczkiewicz A & Orda J. 2002. Digestibility and energy value of non-starch polysaccharides in young chickens, ducks and geese, fed diets containing high amounts of barley. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 131: 657-668. DOI: 10.1016/S1095-6433(01)00517-7
- Jensen MT, Cox RP & Jensen BB. 1995. Microbial production of skatole in the hind gut of pigs given different diets and its relation to skatole deposition in back fat. *Animal Science*, 61: 293-304. DOI: 10.1017/S1357729800013837
- Józefiak D, Rutkowski A, Fratzczak M & Boros D. 2004a. The effect of dietary fibre fractions from different cereals and microbial enzyme supplementation on performance, ileal viscosity and short-chain fatty acid concentrations in the caeca of broiler chickens. *Journal of Animal Feed Sciences*, 13: 487-496. DOI: 10.22358/jafs/67618/2004
- Józefiak D, Rutkowski A & Martin SA. 2004b. Carbohydrate fermentation in the avian ceca: a review. *Animal Feed Science and Technology*, 113: 1-15. DOI: 10.1016/j.anifeedsci.2003.09.007
- Hubener K, Vahjen W & Simon O. 2002. Bacterial responses to different dietary cereal types and xylanase supplementation in the intestine of broiler chicken. *Archiv fur Tierernahrung*, 56: 167-187. DOI: 10.1080/00039420214191
- Komisarczuk S, Merry RJ & McAllan AB. 1987. Effect of different levels of phosphorus on rumen microbial fermentation and synthesis determined using a continuous culture technique. *British Journal of Nutrition*, 57: 279-290. DOI: 10.1079/BJN19870033
- McHan F & Shotts EB. 1993. Effect of short-chain fatty acids on the growth of *Salmonella typhimurium* in an in vitro system. *Avian Disease*, 37: 396-398. DOI: 10.1111/1523-1747.ep12551667
- Metzler-Zebeli BU, Mosenthin R, Baumgärtel T & Rodehutschord M. 2008. The effect of dietary phosphorus and calcium level, phytase supplementation and ileal infusion of pectin on the chemical composition and carbohydrase activity of fecal bacteria and the level of microbial metabolites in the gastrointestinal tract of pigs. *Journal of*

- Animal Science, 86: 1544-1555. DOI: 10.2527/jas.2007-0267
- Metzler-Zebeli BU, Vahjen W, Baumgartel T, Rodehutsord M & Mosenthin R. 2010. Ileal microbiota of growing pigs fed different dietary calcium phosphate levels and phytase content and subjected to ileal pectin infusion. *Journal of Animal Science*, 88: 147-158. DOI: 10.2527/jas.2008-1560
- Miller TL & Wolin MJ. 1974. A serum bottle modification of the Hungate technique for cultivating obligate anaerobes. *Applied Microbiology*, 27: 985-987.
- Minitab. 2010. Minitab 16. Statistical Software. Minitab Inc. State College. Pennsylvania, USA.
- Ptak A, Bedford MR, Swiatkiewicz S, Zyla K & Jozefiak D. 2015. Phytase modulates ileal microbiota and enhances growth performance of the broiler chickens. *PLOS One*, 10: e0119770. DOI: 10.1371/journal.pone.0119770.
- Ricke S. 2003. Perspectives on the use of organic acids and short chain fatty acids as antimicrobials. *Poultry Science*, 82: 632-639. DOI: 10.1093/ps/82.4.632
- Russell JB. 1992. Another explanation for the toxicity of fermentation acids at low pH: anion accumulation versus uncoupling. *Journal of Applied Microbiology*, 73: 363-370. DOI: 10.1111/j.1365-2672.1992.tb04990.x
- Silva SSP & Smithard RR. 2002. Effect of enzyme supplementation of a rye-based diet on xylanase activity in the small intestine of broilers, on intestinal crypt cell proliferation and on nutrient digestibility and growth performance of the birds. *British Poultry Science*, 43: 274-282. DOI: 10.1080/0007166012012150
- Smulikowska S, Czerwiński J & Mieczkowska A. 2010. Effect of an organic acid blend and phytase added to rapeseed cake-containing diet on performance, intestinal morphology, caecal microflora activity and thyroid status of broiler chickens. *Journal of Animal Physiology and Animal Nutrition*, 94: 15-23. DOI: 10.1111/j.1439-0396.2008.00876.x
- van der Wielen PWJJ, Biesterveld S, Notermans S, Hofstra H, Urlings BAP & Van Knapen F. 2000. Role of volatile fatty acids in development of the cecal microflora in broiler chickens during growth. *Applied and Environmental Microbiology*, 66: 2536-2540. DOI: 10.1128/AEM.66.6.2536-2540.2000.Updated
- Widodo AE, O'Neill HM & Iji PA. 2011. Energy utilization and productivity of broiler chickens on diets containing triticale and supplemented with microbial enzymes. *Australasian Medical Journal*, 4: 739.
- Widodo AE, Nolan JV & Iji PA. 2015. The nutritional value of new varieties of high-yielding triticale: Feeding value of triticale for broiler chickens. *South African Journal of Animal Science*, 45: 74-81. DOI: 10.4314/sajas.v45i1.9

# Response of Broiler Chickens to Triticale-Based Diets Supplemented with Microbial Enzymes (2. Microbial Profiles and Activities)

---

## ORIGINALITY REPORT

---

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

18%

PUBLICATIONS

5%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Widodo, AE, JV Nolan, and PA Iji. "The nutritional value of new varieties of high-yielding triticale: Feeding value of triticale for broiler chickens", South African Journal of Animal Science, 2015. 2%

Publication
  - 2** bioone.org 2%

Internet Source
  - 3** D. Józefiak, R.M. Engberg, A. Ptak. "Chapter 2 Relevance of gut microbiome and endogenous phosphatases on phytate breakdown in poultry – implications on phosphorus utilisation", Wageningen Academic Publishers, 2016 1%

Publication
  - 4** M. Akter, H. Graham, P. A. Iji. "Response of broiler chickens to different levels of calcium, non-phytate phosphorus and phytase", British Poultry Science, 2016 1%

Publication
-

5	<a href="http://jwpr.science-line.com">jwpr.science-line.com</a> Internet Source	1%
6	V. García, P. Catalá, J. Madrid, J. Orengo, F. Hernández. "Polysaccharidase preparations added to a wheat-based diet: effects on performance and digestive parameters of broiler chickens held at three different locations", British Poultry Science, 2008 Publication	1%
7	Y. Yang, P. A. Iji, A. Kocher, L. L. Mikkelsen, M. Choct. "Effects of Xylanase on Growth and Gut Development of Broiler Chickens Given a Wheat-based Diet", Asian-Australasian Journal of Animal Sciences, 2008 Publication	1%
8	<a href="http://tandfonline.com">tandfonline.com</a> Internet Source	1%
9	<a href="http://world-food.net">world-food.net</a> Internet Source	1%
10	Ayodeji Oloyo, Adedamola Ojerinde. "Chapter 1 Poultry Housing and Management", IntechOpen, 2020 Publication	1%
11	<a href="http://library.au.dk">library.au.dk</a> Internet Source	1%

Sarbast K. Kheravii, Robert A. Swick, Mingan

12

Choct, Shu-Biao Wu. "Effect of oat hulls as a free choice feeding on broiler performance, short chain fatty acids and microflora under a mild necrotic enteritis challenge", *Animal Nutrition*, 2017

Publication

1%

13

[m.scirp.org](http://m.scirp.org)

Internet Source

1%

Exclude quotes Off

Exclude matches < 50 words

Exclude bibliography Off

# Response of Broiler Chickens to Triticale-Based Diets Supplemented with Microbial Enzymes (2. Microbial Profiles and Activities)

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---