

<http://dx.doi.org/10.29244/jpsl.13.1.88-101>
<http://journal.ipb.ac.id/index.php/jpsl>

About the Journal


Focus and Scope

JPSL publishes articles in fields:

Environmental Policy and Management, Disaster Mitigation, Regional Planning, Land Resources Evaluation, Hidrology, Systems Modelling and Sciences, Water Pollution, Air Pollution, Environmental Technology, Ecotourism, Biodiversity, Environmental Economics, Public Communications, Social Sciences, Anthropology, Environmental Health

Peer Review Process

Submitted papers are evaluated by anonymous referees by double blind peer review for contribution, originality, relevance, and presentation. Editor in Chief will assign the manuscript to a corresponding Section Editor for further handling. The Section Editor will request at least 3 related scientists to review the manuscript. Based on the comments from the Reviewers, Section Editor and Editor in Chief will make the decision on the manuscript.



*click to enlarge

Publication Frequency

Published four times a year: March, June, September, December

Open Access Policy

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

All articles published Open Access will be immediately and permanently free for everyone to read and download. We are continuously working with our author communities to select the best choice of license options, currently being defined for this journal as follows:

- Creative Commons Attribution (CC BY)

How To Register

1. Author should register through the following address: <http://journal.ipb.ac.id/index.php/jpsl/user/register>
2. Author should fulfill the form, especially on the star mark. On the Confirmation Register section, **please give your selection as author, reviewer, or both of them.** After all the form was filled, click on "Register" to process the registration.
3. Please go to User Home and select your role as Author. In Active Submission, click on "Click here: to go to step one of the five-step submission process".
4. These are the five-step submission process:
 - Step 1. Starting the Submission On the Journal Section, select the appropriate section of journal, i.e. Articles. Author has to check-mark on the submission checklists. Author is

able to give brief information to Editor about the submitted manuscript on the Comment form. Please click on the "Save and continue" button to go to the further step.

- Step 2. Entering the Submission's Metadata.
Author should enter metadata of all authors engaged in the writing process. The information of manuscript title, abstract, and keywords must be filled in the textbox.
- Step 3. Uploading the Submission
To upload manuscript to this journal, author should click Browse to find the manuscript file on the hard drive. Choose the file and click Upload button. Once the submission is uploaded, click Save and Continue on the bottom of this page.
- Step 4. Uploading Supplementary Files
The supplementary file should be uploaded is form A, consisting of Statement of Originality and Copyright Release Form. Therefore, click on Browse button, choose the file, click on Upload button, and click on Save and Continue.
- Step 5. Confirming the Submission
Author has to check that all documents are complete. To submit the manuscript to Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan (JPSL), click Finish Submission button. The corresponding author or the principal contact will receive an acknowledgement by email and will be able to view the submission's progress through the editorial process by logging in to the journal web address site.

Author is able to trace the submission manuscript status including manuscript review and editorial process by logging in to the journal web.

Sponsors

- [Pusat Penelitian Lingkungan Hidup-Lembaga Penelitian dan Pengabdian kepada Masyarakat IPB](#)
- [Program Studi Pengelolaan Sumberdaya Alam dan Lingkungan Sekolah Pascasarjana IPB](#)

Journal History

Natural Resources and Environmental Management Studies Program, Graduate School of IPB University (PS-PSL, SPs IPB) in collaboration with Center for Environmental Research - Institute for Research and Community Service, IPB University (PPLH-LPPM, IPB) has been managing JPSL since 2011, with a period of two numbers per year. From 2017 to 2018, JPSL has published three issues per year. Since 2019, JPSL has published four issues per year.

The Journal of Natural Resources and Environment Management (JPSL) is a scientific periodical that presents scientific articles, conceptual thoughts, reviews, and book reviews on natural and environmental resource management that have tropical characteristics. Each manuscript submitted to the Journal of Natural Resources and Environmental Management is reviewed by reviewers.

JPSL is indexed by:

JPSL has been indexed in DOAJ (Directory of Open Access Journal), Google Scholar, Indonesian Publication Index (IPI), and CrossRef. JPSL has also been accredited by Kemenristekdikti as SINTA 2. Valid from January 2022 to January 2026.

Journal Title	: Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan (Journal of Natural Resources and Environmental Management)
Initials	: JPSL
Abbreviation	: Journal of Nat Res EnvMan
Frequency	: 4 (March, June, September, December)
DOI	: 10.29244/jpsl
Print ISSN	: 2086-4639
Online ISSN	: 2460-5824
Editor-in-Chief	: Dr. Yudi Setiawan
Associate Editor	: Dr. Tjahjo Tri Hartono
Publisher	: IPB University, Indonesia
Language	: English
Citation Analysis	: Scopus , Google Scholar
Open Access Policy	: Creative Commons Attribution (CC BY)

Natural Resources and Environmental Management Studies Program, Graduate School of IPB University (PS-PSL, SPs IPB) in collaboration with Center for Environmental Research - Institute for Research and Community Service, IPB University (PPLH-LPPM, IPB) has been managing JPSL since 2011, with a period of two numbers per year. From 2017 to 2018, JPSL has published three issues per year. Since 2019, JPSL has published four issues per year. The Journal of Natural Resources and Environment Management (JPSL) is a scientific periodical that presents scientific articles, conceptual thoughts, reviews, and book reviews on natural and environmental resource management that have tropical characteristics. Each manuscript

submitted to the Journal of Natural Resources and Environmental Management is reviewed by reviewers.

JPSL has been indexed in DOAJ (Directory of Open Access Journal), Google Scholar, Indonesian Publication Index (IPI), and CrossRef. JPSL has also been accredited by Kemenristekdikti as SINTA 2. Valid from January 2022 to January 2026.

Editorial Team

Editor-in-Chief



• **Dr. Yudi Setiawan, SP., M.Env.Sc**



- Scopus ID : [36697084100](https://scopus.com/authid/detail.url?authorID=36697084100)
- ORCID ID <https://orcid.org/0000-0002-1482-1905>
- Google Scholar ID [CF3xiFEAAAAJ](https://scholar.google.com/citations?user=CF3xiFEAAAAJ)
- IPB University, **Indonesia**

Associate Editor



•

Dr. Tjahjo Tri Hartono

- Google Scholar ID [mD3jDtUAAAAJ](#)
- IPB University, **Indonesia**

Section Editor

Dr. Zaenal Abidin, Magr (Scopus ID: [57201126477](#); Google Scholar: [XSdb8AAAAJ&h](#))
*Department of Chemistry, IPB University, **Indonesia***

Dr. Syartinilia, SP., M.Si (Scopus ID: [23668896000](#); Google Scholar: [3YfyIAAAAAJ&h](#))
*Department of Landscape Architecture, IPB University, **Indonesia***

Dr. Perdinan, MNRE (Scopus ID: [56153733100](#); Google Scholar: [bhYh8AAAAJ&h](#))
*Southeast Asian Regional Centre for Tropical Biology, **Indonesia***

Dr. Liyantono, S.TP., M.Agr (Scopus ID: [54906200300](#); Google Scholar: [O52WZvsAAAAJ](#))
*Department of Mechanical and Bio-system Engineering, IPB University, **Indonesia***

Dr. Ir. Sri Mulatsih, M.Sc (Scopus ID: [57202135016](#); Google Scholar: [80SPHIQAAAAJ](#))
*Department of Economics, IPB University, **Indonesia***

Dr. Andrea Emma Pravitasari, SP., Msi (Scopus ID: [56544593100](#); Google Scholar: [le6pzp0AAAAJ](#))
*Department of Soil Science and Land Resources, IPB University, **Indonesia***

Dr. Syamsul Bahri Agus, M.Si (Scopus ID: [56728270900](#); Google Scholar: [jACtQ2IAAAAAJ](#))
*Department of Marine Science and Technology, IPB University, **Indonesia***

Editorial Board

Prof. Mark Cochrane (Scopus ID: [57208455126](#); Google Scholar: [4cu13ecAAAAJ](#))
*Appalachian Laboratory, University of Maryland Center for Environmental Science, **USA***

Dr. Murray Collins (Scopus ID: [37053758600](#); Google Scholar: [QoOxsAAAAJ&h](#))
*School of Geosciences, University of Edinburgh, **United Kingdom***

Prof. Misa Masuda (Scopus ID: [35795906500](#); Google Scholar:)
*Faculty of Life and Environmental Sciences, University of Tsukuba, **Japan***

Prof. Mamoru Kanzaki (Scopus ID: [7006880712](#); Google Scholar:)
*Graduate School of Agriculture, Kyoto University, **Japan***

Prof. Kunihiko Yoshino (Scopus ID: [55611842700](#); Google Scholar:)
Graduate School of Agricultural and Life Sciences, University of Tokyo, Japan

Dr. Takahiro Ota (Scopus ID: [57077829600](#); Google Scholar: [CshOoAAAAJ&h](#))
Graduate School of Fisheries and Environmental Science, Nagasaki University, Japan

Dr. Nishida Kenlo Nasahara (Scopus ID: [24178270500](#); Google Scholar: [xT1dn1IAAAAJ](#))
Faculty of Life and Environmental Sciences, University of Tsukuba, Japan

Dr. Mohamed Kefi (Scopus ID: [36696198900](#); Google Scholar: [QoOxsAAAAJ&h](#))
Water Resource Management, Water Resource and Technology Center (CERTE), Tunisia

Dr. Anis Chekirbane (Scopus ID: [56422279800](#); Google Scholar: [2bqehEUAAAAAJ](#))
Water Resource Management, Water Resource and Technology Center (CERTE), Tunisia

Dr. Tien Dat Pham (Scopus ID: [57188874343](#); Google Scholar: [i9sIr3YAAAAAJ](#))
School of Natural Sciences, Macquire University, Australia

Dr. Gay Jane Perez (Scopus ID: [7203067180](#); Google Scholar: [Kqm_MFUAAAAAJ](#))
University of the Philippines Diliman, Philippines

Prof. Dr. Kasturi Devi Kanniah (Scopus ID: [6503862357](#); Google Scholar: [YFcCk74AAAAAJ](#))
Universiti Teknologi Malaysia, Malaysia

Dr. Rudiyanto (Scopus ID: [57195546540](#); Google Scholar: [4cu13ecAAAAAJ](#))
Universiti Malaysia Terengganu Malaysia

Dr. Muhammad Rokhis Khomarudin (Scopus ID: [55499730500](#); Google Scholar: [RA5YWWQAAAAAJ](#))
Indonesian National Institute of Aeronautics and Space, National Research and Innovation Agency (BRIN), Indonesia

Dr. Dony Kushardono (Scopus ID: [57193562396](#); Google Scholar: [5HnuNO4AAAAAJ](#))
Indonesian National Institute of Aeronautics and Space, National Research and Innovation Agency (BRIN), Indonesia

Dr. Didit Okta Pribadi (Scopus ID: [56433387200](#); Google Scholar: [AiaHhZUAAAAAJ](#))
Indonesia Institute of Science, National Research and Innovation Agency (BRIN), Indonesia

Dr. Luki Subehi (Scopus ID: [25958454100](#); Google Scholar: [MXROp4AAAAAJ](#))
Indonesia Institute of Science, National Research and Innovation Agency (BRIN), Indonesia

Prof. Indayati Lanya (Scopus ID: [57192950522](#); Google Scholar: [RDtpCTYAAAAAJ](#))
Udayana University, Indonesia

Dr. I Wayan Gede Astawa Karang (Scopus ID: [54408470900](#); Google Scholar: [7Rn05yjKDpUC](#))
Udayana University, Indonesia

Dr. Muhammad Kamal (Scopus ID: [55266523600](#); Google Scholar: [J6qAj8kAAAAJ](#))
Gadjah Mada University, Indonesia

Dr. Muhammad Helmi (Scopus ID: [57193740263](#); Google Scholar: [ndG7MCcAAAAJ](#))
Gadjah Mada University, Indonesia

Dr. Joga Dharma Setiawan (Scopus ID: [6603605682](#); Google Scholar: [E2hskOQAAAAJ](#))
Gadjah Mada University, Indonesia

Dr. Elham Sumarga (Scopus ID: [56189783300](#); Google Scholar: [sXh8mGMAAAAJ](#))
Bandung Institute of Technology, Indonesia

Prof. Dr. Lina Karlinasari, S.Hut.MSc.F (Scopus ID: [24171272000](#); Google Scholar: [dDIF1dcAAAAJ](#))
Department of Forestry Product, IPB University, Indonesia

Prof. Dr. Ir. Widiatmaka, DAA (Scopus ID: [56962708400](#); Google Scholar: [jFpFZigAAAAJ](#))
Department of Soil Science and Land Resources, IPB University, Indonesia

Prof. Dr. Ir. Surjono Hadi Sutjahjo, M.S (Scopus ID: [54897314000](#); Google Scholar: [Wf48DbgAAAAJ](#))
Department of Agronomy and Horticulture, IPB University, Indonesia

Prof. Dr. Ir. Sumardjo, M.S (Scopus ID: [56655181500](#); Google Scholar: [zmUmScoAAAAJ](#))
Department of Community Development and Communication Science, IPB University, Indonesia

Prof. Dr. Ir. Lilik Budi Prasetyo, M.Sc (Scopus ID: [35762297500](#); Google Scholar: [wL52oAAAAJ&h](#))
Department of Forest Resources Conservation and Ecotourism, IPB University, Indonesia

Prof. Dr. Ir. Hefni Effendi, M.Sc (Scopus ID: [54922085400](#); Google Scholar: [1g8nkAAAAJ&h](#))
Center for Environmental Research, Institute of Research and Community Services, IPB, Indonesia

Prof. Dr. Ir. Hadi Susilo Arifin, M.S (Scopus ID: [15080540000](#); Google Scholar: [AalJIAAAAJ&h](#))
Department of Landscape Architecture, IPB University, Indonesia

Prof. Dr. Ir. H. Cecep Kusmana, MS (Scopus ID: [30467860100](#); Google Scholar: [T_IbkAAAAJ&h](#))
Department of Silviculture, IPB University, Indonesia

Prof. Dr. Ir. Akhmad Fauzi, M.Sc (Scopus ID: [6602797726](#) ; Google Scholar: [T65DoAAAAJ&h](#))
Department of Resources and Environmental Economics, IPB University, Indonesia

Dr. Ir. Tania June, M.Sc (Scopus ID: [13008836600](#); Google Scholar: [M1WGYAAAAJ&h](#))
Department of Geophysics and Meteorology, IPB University, Indonesia

Dr. Ir. Soeryo Adiwibowo, M.S (Scopus ID: [35087987300](#); Google Scholar: [j2dM0AAAAJ&h](#))
Department of Community Development and Communication Sciences, Indonesia

Dr. Ir. Mohammad Yani, M.Eng (Scopus ID: [6603321602](#); Google Scholar: [coO5AAAAAJ&h](#))
Department of Agroindustrial Technology, IPB University, Indonesia

Dr. Ir. Mirza Dikari Kusri, M.Si (Scopus ID: [16316341900](#); Google Scholar: [nWL0oAAAAJ&h](#))
Department of Forest Resources Conservation and Ecotourism, IPB University, Indonesia

Dr. Ir. Dodik Ridho Nurrochmat, M.Sc.F (Scopus ID: [56403310100](#); Google Scholar: [ITv7pkwAAAAJ](#))
Department of Forests Management, IPB University, Indonesia

Dr. Ir. Arief Sabdo Yuwono, M.Sc (Scopus ID: [56131659400](#); Google Scholar: [_TkeQAAAAJ&h](#))
Department of Civil and Environmental Engineering, IPB University, Indonesia

Dr. Ir. Aceng Hidayat, M.T (Scopus ID: [57200071935](#); Google Scholar: [O-1eVigAAAAJ](#))
Department of Resources and Environmental Economics, IPB University, Indonesia

Dr. I Wayan Nurjaya, M.Sc (Scopus ID: [6505544315](#) ; Google Scholar: [mkWb0U8AAAAJ](#))
Departement of Marine Science and Technology, IPB University, Indonesia

Treasurer



• **Lasriama Siahaan, M.Si**

- IPB University, **Indonesia**

Copyeditor



Luisa Febrina Amalo, S.Si

- Scopus ID : [57193567784](https://scopus.com/authid/detail.url?authorID=57193567784)
- ORCID ID <http://orcid.org/0000-0003-3306-8581>
- IPB University, **Indonesia**

Layout Editor



Vidya Nur Trissanti, S.Hut

- IPB University, **Indonesia**

Proofreader



•

Sylvannisa Putri, S.Hut

- IPB University, **Indonesia**

Website Administrator



•

Arif Kurnia Wijayanto, S.TP, M.Sc

- Scopus ID : [57188697141](https://scopus.org/authorid/57188697141)
- ORCID ID <http://orcid.org/0000-0003-4581-6065>
- Google Scholar ID [xfQppT0AAAAJ](https://scholar.google.com/citations?user=xfQppT0AAAAJ)
- IPB University, **Indonesia**

[Current Issue](#)

Current Issue

Vol. 13 No. 1 (2023): Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan (JPSL)

Published: 2023-03-11

Articles

- **[Spectroscopic Analysis and Dynamics of Dissolved Organic Carbon from an Oil Palm Plantation Peatland](#)**

Untung Sudadi, Ahmad Imtaz Sumbari, Budi Nugroho

1-10

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.1-10](https://doi.org/10.29244/jpsl.13.1.1-10)

- • **[Analysis of Potential, Attraction, and Tourist Perception of Limpakuwus Pine Forest Natural Tourism, Banyumas City](#)**

Alya Utami, Achmad Iqbal, Rahab Rahab

11-26

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.11-26](https://doi.org/10.29244/jpsl.13.1.11-26)

- • **[Phytoplankton response to differences in light and the addition of phosphorus in Lake Cikaret](#)**

Fachmijany Sulawesty, Sata Yoshida Sri Rahayu, Yustiwati, Tjandra Chrismada

27-36

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.27-36](https://doi.org/10.29244/jpsl.13.1.27-36)

- • **[SPECIES COMPOSITION, STRUCTURE, AND MANAGEMENT IN GAYO COFFEE-BASED AGROFORESTRY SYSTEM](#)**

[The Case of Mude Nosar Village, Central Aceh Regency](#)

Adisti Permatasari Putri Hartoyo, Labana Hutagalung, Fifit Kulsum, Arzyana Sunkar, Elis Nina...

37-48

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.37-48](https://doi.org/10.29244/jpsl.13.1.37-48)

• • **[Economic Valuation of Sun Bear \(*Helarctos malayanus*\) and Sumatran Elephant \(*Elephas maximus sumatranus*\) Based on Maintenance Cost Approach](#)**

Audelia Thalita Ramadhanti, Yanto Santosa, Arzyana Sunkar

49-56

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.49-56](https://doi.org/10.29244/jpsl.13.1.49-56)

• • **[Modeling The Habitat Suitability of Dare Monkey \(*Macaca maura* Hr Schinz 1825\) in Bantimurung Bulusaraung National Park, South Sulawesi](#)**

gede tragya suartika pramana, Lilik Budi Prasetyo, Entang Iskandar

57-67

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.57-67](https://doi.org/10.29244/jpsl.13.1.57-67)

• • **[Resource Equivalency Analysis \(REA\): Implication environmental of groundwater in Kupang East Nusa Tenggara](#)**

Aplonia Bani, Suprihatin, SK Saptomo, RL Kaswanto

68-75

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.68-75](https://doi.org/10.29244/jpsl.13.1.68-75)

• • **[THE UTILIZATION OF DURIAN PEELS \(*Durio zibethinus*\) FOR THE MANUFACTURING OF CHARCOAL BRIQUETTES AS ALTERNATIVE FUEL](#)**

Sitti Rahmawati, Rabasia, Afadil, Suherman, Tri Santoso, Vivi Diah Sangkota, Paulus Hengky Abram

76-87

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.76-87](https://doi.org/10.29244/jpsl.13.1.76-87)

• • **[Biophysical characteristics of Wosi Watershed area in Manokwari Regency, Indonesia](#)**

Mahmud, Abdul Aziz, Danang Wijaya, Wahyudi, Bambang Nugroho, Denisa Melanesia

88-101

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.88-101](https://doi.org/10.29244/jpsl.13.1.88-101)

• • **[Challenges and opportunities: application of clean and affordable energy in the development of the national capital of the archipelago](#)**

Dodi Al Vayed, Eko Priyo Purnomo

102-113

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.102-113](https://doi.org/10.29244/jpsl.13.1.102-113)

• • **[Economic Impact of the Eel. Eel \(Anguilla Bicolor\) Aquaculture on Local Communities in Banyuwangi Regency](#)**

Osmaleli Osmaleli

114-121

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.114-121](https://doi.org/10.29244/jpsl.13.1.114-121)

• • **[The role of adiwiyata school in the change of students' knowledge, attitude, and behavior towards the environment](#)**

Siti Badriyah Rushayati, Rachmad Hermawan, Lin Nuriah Ginoga

122-128

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.122-128](https://doi.org/10.29244/jpsl.13.1.122-128)

• • **Green awareness of female consumers towards sustainable products in Indonesia**

Wayan Weda Asmara Dewi, Wifka Rahma Syauki

129-139

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.129-139](https://doi.org/10.29244/jpsl.13.1.129-139)

• • **Study of settlements carrying capacity in Jebres Subdistrict, Surakarta with das sein – das sollen approach**

Desta Eka Fahrurozi, Berlian Warit Amalia, Lia Kusumaningrum, Desma Asty Pramudita, Gavriel Enos...

140-146

Abstract View : 0

Download :0

[10.29244/jpsl.13.1.140-146](https://doi.org/10.29244/jpsl.13.1.140-146)



Biophysical characteristics of Wosi Watershed area in Manokwari Regency, Indonesia

Mahmud^a, Abdul Azis^a, Danang Wijaya^b, Wahyudi^a, Bambang Nugroho^a, Denisa Melanesia^a

^a Faculty of Forestry, Papua University, Amban, 98314, Indonesia

^b Watershed Management Department and Protected Forest Remu Ransiki, Manokwari Papua Barat, Indonesia, 98314

Article Info:

Received: 15 - 08 - 2022

Accepted: 24 - 09 - 2022

Keywords:

Manokwari, mitigation, morphometry, river flow, triangular

Corresponding Author:

Mahmud

Faculty of Forestry, Papua University;

Phone: +6285292837916

Email:

mahmudalya6@gmail.com

Abstract. Flood is number one Indonesian natural disaster in the last 10 years and its occurrence at Manokwari is frequently reported. Biophysical condition is playing a key role in carrying capacity of this catchment area. This study is to determine biophysical characteristics of Wosi Watershed to manage and mitigate flooding in Manokwari. Spatial analysis and field observation methods were used to collect the data. Biophysical variables are rainfall, watershed morphometric, slope, and land used. Carrying capacity is measured using flow regime coefficient and annual flow coefficient. The results showed that the heavy rainfall (> 100 mm) throughout the ten years with 10.5 wet months at average resulting very wet tropical climate. This watershed has an area of 2,346.32 ha, its circumference of 29.95 km² with river length of 8.38 km resulting 0.33 (triangle) and 1.027 (triangle) for Rc and Re, respectively. This morphometry is rectangular and slightly oval (triangular) formed of four rivers with drainage pattern of dendritic, which resembles the shape of a tree branch/twig. Steep slopes are dominant (58.5%), with non-forest area (62%) of the flat and steep slope for settlement (698 ha), and flat slope for mixed dry farming (707 ha). From 2016-2020, river water flow changes rapidly from low to very high to generate flooding, but the carrying capacity is sometime changeable from good to bad. Water drainage, retaining walls, replantation, early warning system, and flooding leaflets mitigation campaign, are structural and non-structural mitigation could be parallelly conducted to manage and mitigate the flooding risks in future.

How to cite (CSE Style 8th Edition):

Mahmud, Azis A, Wijaya D, Wahyudi, B, Melanesia D. 2023. Biophysical characteristics of Wosi Watershed area in Manokwari Regency, Indonesia. *JPSL* 13(1): 88–101. <http://dx.doi.org/10.29244/jpsl.13.1.88–101>.

INTRODUCTION

Throughout 2020, there were 2,952 natural disasters in Indonesia. Thirty six percent of these disasters are floods (CNN Indonesia 2020). The trend of flood continues to increase each year, including throughout 2021 and the first semester of 2022. Flood indicates the inability of the watershed ecosystem to provide environmental services as a result of the degradation and deforestation of forest and land resources (Papaioannou et al. 2015). Flood is caused by climate change, forest conversion, and land use. Thus, these factors increase surface runoff. The amount of water that directly becomes runoff increases significantly, so peak discharge becomes larger (Yan et al. 2013; Nasir et al. 2017). The conversion of forests into

plantations, settlements area, and other built-up areas is sometimes not realized. Whereas in general, the rain that occurs in an area does not experience changes that have an impact on stagnant water.

The watershed is part of a complex ecosystem because it involves various bio-geophysical, social, economic, cultural, and institutional components that interact with each other. Interactions between biotic and abiotic components in watershed ecosystems have direct or indirect impacts and large or small effects. According to Junaidi and Tarigan (2011), land use influences the function of a watershed ecosystem. In addition, Supangat (2012) stated that watershed characteristics and land use are two important factors that affect the hydrological characteristics of the watershed. A comprehensive understanding of the hydrological characteristics of the watershed and the influencing factors is needed for sustainable watershed management, particularly with regard to the use of water resources and land use (Azizah et al. 2021).

Watershed characteristics (typology) are vital for watershed management plans based on Government Regulation Number 37, in 2012 concerning Watershed Management. The typology of the watershed can be identified through the analysis of the characteristics of the watershed. Among the characteristics of a watershed that often causes natural disasters, namely, flooding, one of the conditions of water management is resulting from the output of a watershed (Paimin et al. 2012). To determine the characteristics of the watershed is the biophysical condition of the watershed, which is characterized by parameters related to climate, morphometrics, topography, soil, geology, vegetation, land use, hydrology, and humans.

These climatic and biophysical characteristics will provide specific responses from the watershed to rainfall which will affect the character parameters and special features of water systems such as evaporation, transpiration, surface runoff, infiltration, interception, groundwater, and river discharge (Ningkeula 2016). Increased surface runoff has an impact on river discharge which, if the water cannot be accommodated, then flooding is inevitable. These parameters have been widely used in various geo-hydro-morphological studies, such as flood characteristics, sedimentation, and changes in watershed morphology. In addition, land use by human activities and the morphology of the watershed also affect the water system, including sedimentation.

Wosi watershed is part of 2,145 watersheds spread across Indonesia whose status “needs to be restored” (KLHK 2019). The Wosi watershed is also part of seven watersheds in Manokwari Regency, which are classified as “critical”. Pamuji and Hardianti (2019) argued that the Wosi River could not accommodate the peak flow during high-intensity of rainfall, thus causing flooding. The Wosi area is located in the capital city of West Papua Province, which will continue to grow and develop, but its carrying capacity is decreasing (Mahmud et al. 2021a). The Wosi watershed has become a source of catastrophic flooding for several regions in Manokwari, particularly those living downstream of the watershed, such as the area of Wosi market, the green valley, around the Rendani Airport, the Bugis Village, and the Javanese Village. This study aims to determine the biophysical condition or characteristics of the Wosi Watershed to examine the biophysical variables that contribute to the water system or hydrology. This information is expected to contribute to the management of the Wosi Watershed and flood mitigation in Manokwari.

METHODS

Research Location

This research was carried out for 4 months in the Wosi Watershed, which is geographically located at 133° 0' 0.722" E – 134° 3' 34.55"E and 0° 50' 25.5" S – 0° 54' 8.24" S with an area of 2,346.32 ha (Figure 1). Downstream of the Wosi Watershed is a built-up area and is the capital of West Papua Province. The upstream of the Wosi Watershed covers a fairly steep hill, and its estuary is in Doreri Bay.




Figure 1 Study site

Data Collection

The research method used was spatial analysis and field observation. Watershed morphology includes: land use and topography observed directly and through digital elevation models. The morphometry includes: the shape of the watershed, the area of the watershed, the level of branching, the density of the flow, the flow pattern, and the length of the main river, which are measured digitally based on the guidelines for identifying characteristics of the watershed. Meanwhile, for the hydrology, the observed data installed on the *Stasiun Pengamatan Arus Sungai* (SPAS) consists of flow regime coefficients (FRC) and annual flow coefficients (AFC). Rainfall data in the last 10 years obtained from the BMKG Rendani, Manokwari Regency include: daily, monthly, and annual rainfall, wet months, dry months, and rainfall intensity. Wet month (WM): the amount of rainfall is more than 100 mm/month, humid month (HM): the amount of rainfall is between 60 - 100 mm/month, and dry month (DM): the amount of rainfall is less than 60 mm/month. The rainfall data is used to classify the climate according to Schmidt and Fergusson (1954) in (Soewarno 1991).

$$Q = \frac{DM}{WM} \times 100\%$$

Note: Q = climate; DM = number of dry months; WM = number of wet months

To determine the shape of the watershed, it is necessary to value the circulation ratio or RC (Strahler 1964) and the extension ratio or RE (Schum 1956) with the following formula:

$$RC = \frac{4\pi A}{P^2}$$

$$RE = 1,129 \left\{ \frac{A^{0.5}}{L} \right\}$$

Note: RC = circulation ratio; RE = elongation ratio; A = watershed area (m²); L = length of the watershed; P = perimeter of the watershed (m); = 3.14 If RE < RC means the watershed is rounded and RE > RC means the watershed is elongated. Data on circulation ratio, extension ratio, total river length, and watershed circumference were obtained through a geographic information system (GIS). River density number of long rivers, including tributaries and wide watersheds, was obtained through GIS. River density is an index that shows the number of rivers and tributaries in a watershed. The index is obtained by the following equation (Soewarno 1991):

$$Dd = \frac{L}{A}$$

Note: Dd = river density; L = total length of the river, including its tributaries; A = watershed width. The flow regime coefficient (FRC) is obtained from the comparison between the maximum discharge (Q_{max}) and the minimum discharge (Q_{min}) in a watershed, referring to the regulation of minister Number 61/Menhut-II/2014.

$$FRC = \frac{Q_{max}}{Q_{min}}$$

FRC ≤ 20 = very low; 20 < FRC ≤ 50 = low; 50 < FRC ≤ 80 = moderate; 80 < FRC ≤ 110 = high; FRC > 110 = very high. The annual flow coefficient (AFC) is obtained from the comparison between the annual flow rate (Q) and the annual rainfall thickness (P) with the following calculation:

$$AFC = \frac{Q}{P}$$

P is obtained from the annual rainfall x the area of the Wosi watershed. AFC ≤ 0.2 = very low; 0.2 < AFC ≤ 0.3 = low; 0.3 < AFC ≤ 0.4 = moderate; 0.4 < AFC ≤ 0.5 = high; AFC > 0.5 = very high (Permen N0.61/Menhut-II/2014).

Data analysis

Meteorological, morphological, and morphometric data were interpreted to obtain a descriptive qualitative description of the Wosi Watershed characteristics using the ArcGIS 10.5 application. Analysis for hydrological impacts begins with determining the value, weight, and score of the AFC and FRC indicators which refer to Regulation of Minister N0. 61/Menhut-II/2014.

RESULTS AND DISCUSSION

Watershed characteristics show specific characteristics related to water systems, such as soil type, land use, topography, slope, and slope length. Evaporation, transpiration, infiltration, surface runoff, soil water content, and water discharge are greatly dependent on the characters of the watershed in response to rainwater, which are rainfall, watershed shape, river density, watershed slope, and land use.

Rainfall

One of the sources of water on the earth's surface comes from rainfall. When it rains, before the water is absorbed by the soil in the vegetated area, the water will be temporarily held in the vegetation crown and then flow through the stem and the canopy. If a high intensity of rain occurs, then the soil cannot absorb

water then the water will stagnate or flow on the ground surface. The surface runoff will go to lower areas, such as water bodies such as reservoirs, lakes, and rivers. High intensity of rainfall can threaten water quality, causing landslides, sedimentation, and even flooding (Vannier 2016). According to Biswas et al. (2017), the distribution and amount of rainfall greatly affect the occurrence of floods. Rainwater that reaches the ground can become surface runoff and infiltrates the ground (Asdak 2010; Ngongondo et al. 2011; Zhang et al. 2017). Low-absorbed rainwater will cause water to stagnate. Annual rainfall data is grouped based on the criteria of the wet month, medium month, and dry month to get the type of climate in the Wosi Watershed area (Table 1).

Based on the Schmidt and Fergusson system, the Wosi Watershed has a very wet tropical climate type/climate type A (0.028) with a value of $Q = 0 < Q < 0.143$. As a result of the very wet tropical climate/climate type A in the Wosi Watershed and its surroundings, it receives sunlight throughout the year, and air pressure in areas with tropical climates is relatively low and changes regularly and slowly. Evaporation through the soil, vegetation, and water bodies is high. Thus, there are many clouds. The clouds gather together, and a large amount of water droplets cannot be held in the atmosphere. Moreover, soils in tropical climate areas are relatively more fertile as a result of the high intensity of rainfall. In addition, the difference in temperature is not significant between day and night.

Table 1 Wet month, medium month, and dry month

Year	Dry month (< 60 mm)	Medium month (60 - 100 mm)	Wet month (> 100 mm)
2012	0	0	12
2013	1	1	10
2014	0	1	11
2015	1	0	11
2016	0	0	12
2017	0	1	11
2018	0	2	10
2019	1	3	8
2020	0	3	10
2021	0	2	10
Amount	3	13	105
Average	0.3	1.3	10.5

According to Zhang et al. (2017); Worman et al. (2017); Fuchs et al. (2016), the relatively large runoff is caused by high rainfall, and the presence of vegetation is low. Table 1 shows that in the last ten years, it has been dominated by rain/wet months (>100 mm). The impact of heavy rainfall on the damaged area causes greater surface runoff. If the capacity of the river is full, the water will overflow on the ground and into settlements.

Water in the forest, litter, or soil forms clouds through the process of evapotranspiration. It is a cycle that creates a good pattern of rainfall so that long droughts and the danger of flooding will be avoided. The tree has different evapotranspiration and infiltration capabilities based on the species so that surface runoff and erosion can be reduced (Budirianto 2013). The river bank that should have been trees has turned into a plant that can't reduce runoff and sediment that enters the river. Trees with deep roots and dense crowns have turned into fibrous plants that are uprooted and washed away if there is a heavy runoff.

Vannier (2016) argued that flood often occurs if the rainfall lasts long, the distribution is uneven, and the river capacity is limited. For example, in 2017, the Cimanuk flood was caused, among others, by high rainfall (110 to 255 mm/day), land use that was not in accordance with its capabilities, and forest area, which was only 17.9% (Savitri and Pramono 2017). However, according to Mahmud et al. (2018), rainfall is not the

only cause of flooding. Flood is caused by land conversion, sedimentation, river narrowing, and watershed damage. Floods will not occur if the land is able to absorb large amounts of water, the river flows smoothly, the water is accommodated in the river, and the community does not reduce the cross-sectional area of the river. On the other hand, according to Neuvel and Knaap (2010), flooding will occur if the rain is not absorbed by the soil, the water is not accommodated in water bodies, the narrowing of the river has an impact on the flow of water is not smooth, and the community reduces the size of the river bank.

Watershed Morphometric

Watershed morphometric is a term used to express the river channel, such as the area, length, width, slope, river branching level, river density, and the shape of the watershed. The shape of the watershed has an important meaning in relation to river flow/flow velocity. The shape of the watershed allegedly affects the water to reach the outlet (the endpoint of the water release). The morphometric of the Wosi Watershed is shown in Table 2.

Table 2 Morphometric conditions of the Wosi Watershed

Wosi watershed	Watershed morphometry	Wosi watershed	Watershed morphometry
Watershed Area (A)	23,463,200 m ²	Watershed Center Point (Cg)	X = 4.5; Y = 4.08
Main River Length (Lb)	8.38 km	River Flow Mean Slope (So)	0.79%
Watershed Width (W)	3.01 km	Circulation Ratio (Rc)	0.33
Watershed Length (Ln)	5.32 km	Limniscate Constant (k)	1.01
Density River Flow (Dd)	0.35 Height	Biforcation Ratio (Rb)	Rb ^{1/2} : 3; Rb ^{2/3} :
Flow Length (Lg)	4.63 km	Elongation ratio (Re)	2
Watershed Average Elevation (Ma)	250 m asl	Watershed Average Slope (Sb)	1.027
Circumference of the watershed (p)	29.95 km	Weighted average (Wrb)	0.126
Longest River (L)	8.38 km	Factor Symmetry (SIM)	3.75
Coefficient Watershed Form (F)	0.37		1.21

Source: MoLEF 2019

Table 2 shows that the Wosi watershed has an area of 2,346.32 ha, a watershed circumference of 29.95 km², and a river length of 8.38 km, so Rc is 0.33 (triangle) and Re is 1.027 (triangle). From these results, it can be concluded that the shape of the watershed is rectangular and slightly oval (triangular). Paimin et al. (2012) stated that the speed of water flowing to the outlet is influenced by the shape of the watershed. The shorter the time required for water to reach the outlet, the more round the shape of the watershed, so the greater the potential for flooding. On the other hand, the longer the water goes to the outlet if the shape of the watershed is more oval, the flood has low runoff. Likewise, according to Wirosoedarmo et al. (2010), the peak flood discharge value is relatively small, with a relatively long flood peak time in the form of an elongated watershed.




Figure 2 Watershed shape, density, and river network

There are four important rivers flowing in the Wosi Watershed, namely Rendani 1, Rendani 2, Kentek, and Dingin River (Figure 2). The drainage pattern of the Wosi Watershed in general, resembles the shape of a tree branch/twig (dendritic pattern). The river currents in the upstream and middle watersheds are relatively heavy, given the slope of the riverbed is very steep and a massive area of conversion for settlements. The upstream area where the water flows is quite heavy must be maintained as a forest because if there is a conversion to the built-up area, it will easily become a source of erosion. According to Barokah and Purwantoro (2014) upstream areas such as mountainous areas, hills, or mountain slopes are sources of erosion.

Watershed Slope

Slope/topography is the appearance of the land surface caused by the difference in height between two areas. Slope is one of the elements of the occurrence of floods, erosion, and landslides. The steeper and longer the slope, the greater the rate and amount of runoff for major erosion or landslides. On land with a large slope, the surface flow has a large velocity, so infiltration tends to decrease (Triatmodjo 2010; Helman et al. 2017). Soil slope has a very significant influence on soil degradation. An area that is flat and tends to be sunken will be prone to flooding, considering that in this area, water is easily concentrated towards lower areas. The slopes of the Wosi Watershed area are listed in Table 3.

Table 3 shows that there are only two of the slope classes in the Wosi Watershed, which is dominantly steep. The slopes have an influence on the velocity and the volume of surface runoff. Flat topography (0 - 8%) covering an area of 970.58 hectares (41.52%) has become an urban area of West Papua Province,

including Wosi Village, Bugis Village, Javanese Village, Makassar Village, and Rendani. The flat topography of the lower and middle watersheds has become overcrowded into built-up areas and has begun to shift towards steeper slopes.

Table 3 Slope class in the Wosi Watershed

Slope (%)	Area (ha)	Percentage (%)
0 - 8%	970.58	41.52
25 - 40%	1,366.91	58.48
Total	2,337.49	100

According to Oktivova and Rudiarto (2019), the community began to change their behavior in land use in the suburbs from non-built land to built-up land. Furthermore, the steep slopes (25 - 40%) covering an area of 1,366.91 hectares (58.47%) of the Wosi watershed area include the villages of Udopi, Ingramui, and Soribo, with slopes ranging from flat to very steep at an altitude of 0 – 1,450 m from sea level. The steeper the slope, the greater the runoff velocity if the area is less vegetation. Mahmud et al. (2021b) argue that land use for settlements/built-up areas has higher runoff and sediment than forest areas. The results of the survey of areas located on steep slopes that were previously plantations and forest areas began to be converted to settlements (Figure 3).



Figure 3 Ingramui Village settlement adjacent to a steep hill without vegetation (left), a residential area adjacent to a steep hill with vegetation (right)

Figure 3 shows a house building adjacent to a steep slope. The threat of landslides is high. Considering the area has a rather steep and very steep topography, when it rains, the water easily flows on the soil surface. It erodes the top layer of soil and causes the soil to shift/move in large numbers because nothing is able to bind the soil. On the other hand, the area with a rather steep and very steep topography, if it is well maintained (designated as protected forest/protected areas), will maintain soil fertility, protect the soil from the threat of landslides, and provide water, particularly during the dry season.

Hardiyatmo (2012) discovered that topography is very influential on rainfall. If it rains on a steep surface without vegetation, a runoff will arise and flow downstream or to a lower place. Excessive flow cannot be accommodated by the river, causing flooding. In areas with a flat topography to slightly concave, water is easily concentrated and inundated, and if the amount of water is greater, then flood is inevitable (Marfai 2011). One of the flooded locations has built an embankment (Figure 4), but the embankment is not fully able to cope with flooding. Many embankments were built in flooded areas, but because the upstream watershed had a lot of land conversion, water with great speed was able to destroy the embankment, eventually, it broke, and the flood reoccurred.



Figure 4 The flooded area where the embankment has been built

Land Use

The increasing crowded urban areas have led to an increase in land use for residential, agricultural, and water needs. Types of land use around Wosi watershed are divided into 7 groups which are dominated by settlements, as shown in Table 4.

Table 4 Topography and Land Use Functions in the Wosi Watershed

Topography	Use land	Non-forest area (ha)	Forest area (ha)	Percentage (%)	
				Non-forest	Forest
Steep	Primary dryland forest	-	83.15	-	3.54
Steep	Secondary dryland forest	-	773.26	-	32.95
Steep	Bushes /scrub	-	27.64	-	1.17
Flat and Steep	Settlement	697.90	-	29.74	-
Flat and Steep	Open land	3.94	-	0.16	-
Flat	Mixed dryland farming	707.07	-	30.13	-
Flat	Airport	53.35	-	2.27	-
Total		1,462.26	884.05	62.32	37.68

Source: BPDASHL Remu Ransiki 2019

The type of land use as forest based on Table 4 is still above 30%. It is suitable to protect and maintain watershed is at least 30% based on law Number 41 of 1999. The forest cover includes primary dry land forest, secondary dry land forest, and shrubs/shrubs covering an area of 884.05 hectares (37.68%), while the non-forest area is 1,462.26 hectares (62.32%). The forest area of 37.68% is expected to be able to protect and maintain the diversity of flora and fauna, bio-geophysics, and the carrying capacity of natural resources.

Along with the development of Manokwari City as the capital of West Papua Province, the area continues to experience forest land conversion. As Figure 5, the upper Wosi Watershed has been cut off. Some areas are still shaped like a hill where landslides may occur. In fact, in 2012, some of the cities of Manokwari, which are adjacent to the watershed, experienced major floods. City development that does not follow the regency's spatial plan will cause not only a major flood but also landslides. The conversion of land to built-up land is around 28.02 ha (39.5%), resulting in a decrease in groundwater, landslides, and an increase in surface water discharge (Dewi and Rudiarto 2014).



Figure 5 The planned upstream watershed for the built-up area has been partially leveled (left). Some of the lands is still steep and made like a ladder (right)

Impact of Watershed Characteristics on Hydrology

Mahmud et al. (2021a) found out that the carrying capacity of the Wosi Watershed is categorized as bad and is one of the watersheds in Manokwari Regency that must be restored. Water discharge is a very important part of hydrological parameters to evaluate the extent to which rainfall affects surface runoff, river water level, and river capacity. Meanwhile, the FRC is an inseparable part of the water discharge, as shown in Table 5.

Table 5 FRC and AFC values in the Wosi watershed

Year	Max of water discharge (m ³ s ⁻¹)	Min of water discharge (m ³ s ⁻¹)	FRC	Evaluation	Total water discharge (m ³ year ⁻¹)	Total rainfall in the watershed (mm year ⁻¹)	AFC	Evaluation
2016	1.23	0.06	20.53	low	13,038,081.4	48,044,540.9	0.27	low
2017	2.37	0.04	60.57	moderate	12,909,855.2	53,176,793.8	0.24	low
2018	3.73	0.02	232.29	very high	30,749,926.9	33,162,321.5	0.93	very high
2019	7.1	0.01	710	very high	16,018,933.6	67,450,857	0.23	low
2020	6.98	0.02	349	very high	15,986,897.4	66,630,438	0.23	low

Table 5 shows the trend of increasing differences in water flow. The value of FRC from 2016 to 2020 is dominated by very high. This shows that there is a decrease in the carrying capacity of the watershed to receive, absorb, and drain rainwater into the river. According to Mahmud et al. (2021a), if the FRC is low, the water flows throughout the year without showing a significant water level. The flow of water is not only in the rainy season but continues to flow even during the dry season. On the other hand, if the FRC is very high, it means that the flow of river water is easy to change, sometimes low or even very high, which can cause flooding. This indicates that the river flow is bad, because the ability of forests and land to absorb, store, receive rainwater, and release water is bad. Rain intensity, vegetation cover, soil type, slope, and land management techniques are related to the ability of the watershed to receive, store, discharge, and drain water.

River flow is an important part of determining the extent to which surface runoff, river water level, river capacity, and sediment affect rainwater input/output. The annual flow coefficient indicates what percentage of rainfall is a runoff in the watershed. Table 5 also shows that AFC always changes starting from low and very high but is dominated by low. A low AFC means that the watershed's carrying capacity is good, while a very high AFC indicates that the watershed's carrying capacity has decreased.

The increasing rainfall is not always accompanied by an increase in water discharge if there is low rainfall intensity, even distribution of rain, and good canopy cover. More crowns and a wider distribution will reduce the canopy runoff and the flow of stems to reach the soil surface (Mahmud et al. 2019a). When it reaches the ground surface, the water will seep, and if it is saturated, it will become surface runoff. The better the water flow, this indicates that more water flows from the water springs than from the surface runoff. The government and environmentalists must be aware of the community land use to always implement soil and water conservation and follow the spatial plans of Manokwari Regency.

Flooding Management and Mitigation

Recently, there have been massive land use changes in Manokwari Regency for human settlement, local and central government official administration, village development programs, national infrastructure program of a seaport, airport, and roads, industrial expansion, Mahmud et al. (2019a). Forest cover areas conversion into palm oil plantations (Mahmud et al. 2019a), Maruni's forest cover area into cement industries (Mahmud et al. 2020), and Wosi protected forest area into a settlement, and commercial sectors (Mahmud et al. 2021b) have been influenced the hydrological system of Manokwari City and regency. These land use changes supported by the biogeographical characteristic recorded from this research are supposed to generate the occurrence of flooding in Manokwari Regency and town.

Two types of flooding management and mitigation worldwide acknowledge for the tropical area are structural and non-structural systems or measures (Samu and Kentel 2018). Urbanus et al. (2021) stated that management for flooding could be divided into actions, prevention/mitigation, response/intervention, and recovery. Structural mitigation emphasizes the structural and physical in-stream and off-stream such as building water dams, checking dams, ground sill, river normalization, river pass, and others. Non-structural mitigation ranges from law and regulation enforcement, land use policy, replantation, watershed management, public and community awareness, campaign, and education.

Structural and non-structural mitigation could be conducted parallelly to reduce the flooding risk in Manokwari Regency, ranging from making water sanitation/drainage, waterways, and sewer water at the settlement areas, and building retaining walls, gravity walls for the area with a steep slope. Replantation, warning system, flooding leaflets mitigation campaigns, and community involvement are various non-structural mitigation.

CONCLUSION

The Wosi watershed has a very wet tropical climate type/climate type A (0.028) which is dominated by rain/wet months (>100 mm). Its morphometric is a triangular-shaped watershed with 4 rivers, namely Rendani 1 River, Rendani 2 River, Kentek River, and Kali Dingin River. Furthermore, this watershed has flat slopes covering an area of 970.58 hectares and steep slopes occupying an area of 1,366,91 hectares. Land use is dominated by non-forest, with an area of 1,462.26 hectares (62.32%). Meanwhile, the coefficient of flow regime (FRC) is categorized as low, medium, and dominated by very high. Very high FRC means that river water flow changes quickly, sometimes low or even very high, which can cause flooding. The result of the assessment of AFC is categorized as low and very high. A low AFC value means that the carrying capacity of the watershed is good, while a very high AFC indicates that the carrying capacity of the watershed has decreased.

REFERENCES

- [BPDASHL Remu Ransiki] Balai Pengelolaan Daerah Aliran Sungai dan Hutan Lindung Remu Ransiki. 2019. *Laporan Monitoring dan Evaluasi Pengelolaan DAS Wosi Tahun 2019*. Jakarta: Kementerian Lingkungan Hidup dan Kehutanan Republik Indonesia.
- [KLHK] Kementerian Lingkungan Hidup dan Kehutanan. 2019. KLHK dan Kementerian ESDM berkomitmen percepat upaya Reklamasi Hutan dan Rehabilitasi DAS. [accessed 2020 Feb 8]. <http://ppid.menlhk.go.id/berita/siaran-pers/4845/klhk-dan-kementerian-esdm-berkomitmen-Percepat-upaya-reklamasi-hutan-dan-rehabilitasi-das>.
- [Permen] Peraturan Menteri Kehutanan. 2014. Keputusan Menteri Kehutanan No. P. 61/Menhut-II/2014 tentang monitoring dan evaluasi Daerah Aliran Sungai. Jakarta: Kementerian Kehutanan Republik Indonesia.
- [MoF] Ministry of Forestry. 1999. Ministry of Forestry Decree Number 41/Menhut/1999 for Forestry. Jakarta: Ministry of Forestry, Republic of Indonesia.
- [MoLEF] Ministry of Life Environmental and Forestry. 2019. *Laporan Monitoring dan Evaluasi Pengelolaan DAS Arui Tahun 2019*. Jakarta: Ministry of Life Environmental and Forestry.
- Asdak C. 2010. *Hydrology and Management Watershed*. Yogyakarta: Gajah Mada University Press.
- Azizah C, Pawitan H, Nuraida, Satriawan H, Abbas R, Robo S, Misnawati. 2021. Hydrological Characteristics and Its Impact On Flood Jambo Aye Basin In Aceh Indonesia. *JPPDAS*. 5(2):171–184.
- Barokah I, Purwantoro D. 2014. Pengaruh variasi debit aliran terhadap gerusan maksimal di bangunan jembatan dengan menggunakan program HEC-RAS. *INERSIA Jurnal*. 10(2):175–184.
- Biswas S, Kranz WL, Shapiro CA, Snow DD, Hunt SLB, Mamo M, Tarkalson DD, Zhang TC, Shelton DP, Donk SJV, et al. 2017. Effect of rainfall timing and tillage on the transport of steroid hormones in runoff from manure amended row crop fields. *Journal Hazardous Materials*. 324:436–447.
- Budirianto HJ. 2013. Hubungan model arsitektur pohon roux jenis *Koordersiodendron pinnatum* Merr dan koriba Jenis *Pometia pinnata* forster terhadap parameter perimbangan air di Hutan Tanaman Anggori Manokwari. *BIOSCIENTIAE*. 10(1):40–55.
- CNN Indonesia. 2020. *BNPB Ungkap Tren Peningkatan Bencana Banjir hingga Longsor*. [accessed 2021 Feb 03]. <https://www.cnnindonesia.com/nasional/20210202132713-20-601293/bnpb-ungkap-tren-peningkatan-bencana-banjir-hingga-longsor>.
- Dewi NK, Rudiarto I. 2014. Pengaruh Konversi Lahan terhadap Kondisi Lingkungan di Wilayah Peri-urban Kota Semarang (studi kasus: area berkembang Kecamatan Gunungpati). *Jurnal Pembangunan Wilayah dan Kota*. 10(2):115–126.
- Fuchs M, Reverman R, Owen LA, Frankel KL. 2016. Reconstructing the timing of flash floods using dating at Leidy Creek alluvial fan and valley, White Mountains, California–Nevada, USA. *Quaternary Research Journal*. 83(1):178–186.
- Hardiyatmo HC. 2012. *Tanah Longsor dan Erosi: Kejadian dan Penanganan*. Yogyakarta: Gajah Mada University Press.
- Helman D, Osem Y, Yakir D, Lensky IM. 2017. Relationships between climate, topography, water use and productivity in two key mediterranean forest types with different water-use strategies. *Agricultural and Forest Meteorology*. 232:319–330. doi:<https://doi.org/10.1016/j.agrformet.2016.08.018>.
- Junaidi E, Tarigan SD. 2011. Pengaruh hutan dalam pengaturan tata air dan proses sedimentasi daerah aliran sungai (DAS): studi kasus di DAS Cisadane. *Jurnal Penelitian Ilmiah dan Konservasi Alam*. 8(2):155–176.
- Mahmud, Budirianto HJ, Wahyudi, Kusumandari A. 2020. Studi alih fungsi menjadi pertambangan semen pada hutan lindung Maruni Kabupaten Manokwari (A study conversion to be mining cement in Maruni protected forest Manokwari Regency). *Journal of Natural Resources and Environmental Management*. 10(3):545–558. doi:<http://dx.doi.org/10.29244/jpsl.10.3.545-558>.

- Mahmud, Kusumandari A, Sudarmadji, Supriyatno N. 2018. A study of flood causal priority in Arui Watershed, Manokwari Regency, Indonesia. *Jurnal Manajemen Hutan Tropika*. 24(2):81–94.
- Mahmud, Kusumandari A, Sudarmadji, Supriyatno N. 2019b. Conservation design and scenario for flood mitigation on Arui Watershed, Indonesia. *Indonesian Journal of Geography*. 51(3):261–272.
- Mahmud, Kusumandari A, Sudarmadji, Supriyatno N. 2019a. The species diversity and structure of the limited production forest in Arui Watershed of Manokwari District of West Papua, Indonesia. *Biosaintifika Journal of Biology and Biology Education*. 11(2):279–288.
- Mahmud, Wahyudi, Bataradewa S, Budirianto HJ, Mutakim, Muhlis LO. 2021b. The relationship of rainfall on surface runoff and sediments on various land use in Arui Watershed, Manokwari Regency. *Jurnal Ilmu Tanah dan Lingkungan*. 23(2):85–92.
- Mahmud, Wijaya D, Wahyudi, Kusumandari A. 2021a. Evaluation of carrying capacity and conservation scenarios of Wosi Watershed at Manokwari Regency, West Papua. *Jurnal Ilmu Kehutanan*. 15(2):231–246.
- Marfai MA. 2011. Impact of coastal inundation on ecology and agricultural land use case study in Central Java, Indonesia. *Quaestiones Geographicae*. 30(3):19–31.
- Nasir A, Saleh MB, Bahrani 2017. Optimization of land use collaborative management model Perum Perhutani: study case KPH Pekalongan Barat. *Jurnal Manajemen Hutan Tropika*. 23(1):25–36. doi:<https://doi.org/10.7226/jffm.23.1.25>.
- Neuvel JMM, Knaap WVD. 2010. A spatial planning perspective for measures concerning flood risk management. *International Journal of Water Resources Development*. 26(2):283–296. doi:<http://dx.doi.org/10.1080/07900621003655668>.
- Ningkeula ES. 2016. Analisis karakteristik morfometri dan hidrologi sebagai ciri karakteristik biogeofisik DAS Wai Samal Kecamatan Seram Utara Timur Kobi Kabupaten Maluku Tengah. *Jurnal Ilmiah Agribisnis dan Perikanan*. 9(2):76–86.
- Ngongondo C, Xu C, Gottschalk L, Alemaw B. 2011. Evaluation of spatial and temporal characteristics of rainfall in Malawi: A case of data scarce region. *Theoretical and Applied Climatology*. 106:79–94. doi:<https://doi.org/10.1007/s00704-011-0413-0>.
- Oktinova N, Rudiarto I. 2019. Study on land use around bukit Semarang Baru area. *Jurnal Pembangunan Wilayah dan Kota*. 15(4):262–276.
- Paimin, Pramono IB, Purwanto, Indrawati DR. 2012. *Sistem Perencanaan Daerah Aliran Sungai*. Jakarta: Kementerian Kehutanan Republik Indonesia.
- Pamuji KE, Hardianti. 2019. Evaluasi hidrologi Daerah Aliran Sungai Wosi dalam menghadapi curah hujan ekstrim. *Jurnal Natural*. 15(2):67–77.
- Papaioannou G, Vasiliades L, Loukas A. 2015. Multi-criteria analysis framework for potential flood prone areas mapping. *Water Resour Manage*. 29:399–418. doi:<https://doi.org/10.1007/s11269-014-0817-6>.
- Samu R, Kentel AS. 2018. An analysis of the flood management and mitigation measures in Zimbabwe for sustainable future. *International Journal of Disaster Risk Reduction*. 31:691–697. <https://doi.org/10.1016/j.ijdr.2018.07.013>.
- Savitri E, Pramono IB. 2017. Upper Cimanuk flood analysis of 2016. *Journal of Watershed Management Research*. 1(2):97–110.
- Schum SA. 1956. Evolution of drainage system and slopes in badlands at Perth Amboy. *Geological Society of America*. 67(5):597–646.
- Soewarno. 1991. *Hidrologi: Pengukuran dan Pengolahan Data Aliran Sungai (Hidrometri)*. Bandung: Nova.
- Strahler AN. 1964. Quantitative geomorphology of drainage basins and channel networks. In: Chow VT, editor. *Handbook of Applied Hydrology*. New York (NY): McGraw-Hill.
- Supangat AB. 2012. Hydrological characteristics based on morphometric parameters of Meru Betiri National Park. *Jurnal Penelitian Hutan dan Konservasi Alam*. 9(3):275–283.
- Triatmodjo B. 2010. *Hidrologi Terapan*. Yogyakarta: Beta Offset.

- Urbanus A, Sela RLE, Tungka AE. 2021. Mitigasi bencana banjir struktural dan non-struktural di Kabupaten Bolaang Mongondow Selatan. *Jurnal Spasial*. 8(3):447–458.
- Vannier O, Anguetin S, Braud I. 2016. Investigating the role of geology in the hidrological response of mediterranean cathment prone to flash floods: regional modeling study and proses understanding. *Journal of Hydrology*. 541:158–172.
- Wirosoedarmo R, Haji ATS, Pramesti EM. 2010. Study on form, drainage network, and watershed hydrograph by Using SIMODAS (case study on Sabu Island - Nusa Tenggara Timur). *Jurnal Teknologi Pertanian*. 11(2):123–130.
- Worman A, Lindstrom G, Riml J. 2017. The power of runoff. *Journal of Hydrology*. 548:784–793. doi:<https://doi.org/10.1016/j.jhydrol.2017.03.041>.
- Yan B, Fang NF, Zhang PC, Shi ZH. 2013. Impacts of land use change on watershed streamflow and sediment yield: an assessment using hydrologic modelling and partial least squares regression. *Journal of Hydrology*. 494:26–37. doi:<https://doi.org/10.1016/j.jhydrol.2013.01.008>.
- Zhang FB, Bai YJ, Xie LY, Yang MY, Li ZB, Wu XR. 2017. Runoff and soil loss characteristics on loess slopes covered with aeolian sand layers of different thicknesses under simulated rainfall. *Journal of Hydrology*. 549:244–251.