# Rasayan 2021-the influence of soil

by Darma Santi

Submission date: 20-Apr-2023 10:11AM (UTC+0900)

**Submission ID:** 2069843632

File name: Rasayan\_2021-the\_influence\_of\_soil.pdf (1.04M)

Word count: 2394

Character count: 11609



RASĀYAN J. Chem.

Vol. 14 | No. 1 | 454-459 | January - March | 2021 ISSN: 0974-1496 | e-ISSN: 0976-0083 | CODEN: RJCABP http://www.rasayanjournal.com http://www.rasayanjournal.co.in

## THE INFLUENCE OF SOIL CHEMICAL CONTENT FACTOR ON RUSWERI RIVER WATER QUALITY IN SUPIORI REGENCY, PROVINCE OF PAPUA, INDONESIA

J. V. Morin<sup>1,⊠</sup>, B. Mansoben<sup>2</sup>, M. Infaindan<sup>2</sup>, F.L. Ayer<sup>2</sup>, Marty<sup>2</sup>, L. L. Mandosir<sup>2</sup>, J.O. Ughude<sup>2</sup>, D. Awendu<sup>2</sup>, R. Yarangga<sup>2</sup>, P. Patulak<sup>2</sup>, P. Noya<sup>2</sup>, and D. Santi<sup>1,3</sup>

<sup>1</sup>Chemistry Department, Faculty of Mathematics and Nature Sciences Papua University, Papua Barat, Indonesia

<sup>2</sup> Department of Environment Agency, Supiori District, Papua, Indonesia <sup>™</sup> Corresponding Author: jecko morin@yahoo.com

#### ABSTRACT

Research on the effect of chemicals on the water quality of the Rusweri river in Supiori Regency has been conducted. The inventory plan of the river as a provider of raw water in Supiori Regency will begin in 2020, and the Rusweri river is one of the candidates to be used, it will be seen using its green water for microbiological assistance that can be researched the Rusweri river. This study uses a descriptive method that explains the water quality the Rusweri river with physical and chemical monitoring parameter methods for each place in the Rusweri river. The results of this study contain several chemical and physical parameters that exceed the quality standard threshold. Increased concentrations of parameters PO<sub>4</sub>-, Fe, F<sup>-</sup>, COD, and BOD from upstream to downstream due to the river bed structure that already contains particular metal contents and also the influence of organic and microbiological concentrations that enter through the addition of secondary rivers on the body Rusweri river. Calculation of river pollution index results in an average of mildly polluted status, which is dominated by phosphate parameters.

Keywords: Soil quality, Water quality, Rusweri, Papua

23 RASĀYAN *J. Chem.*, Vol. 14, No.1, 2021

#### INTRODUCTION

Water quality an essential role for all living things. Surface water quality in an area is governed by both natural processes such as rainfall levels, weathering and soil erosion processes and anthropogenic effects chas urban, industrial and agricultural activities and human exploitation of water resources. Groundwater quality has become an important water resource problem due to rapid population growth, rapid industrialization, unplanned urbanization, pollution from highlands to lowlands, and overuse of fertilizers, 19 ticides in agriculture 1.2.

Based on the Republic of Indonesia Government Regulations (quality standard) (PP RI) No. 82 of 2001 concerning Management of Water Quality and Control of Water Pollution (river/surface water) and PERMENKES No. 7 of 2019 concerning hospital environmental health, water and soil quality monitoring needs to be done in each District/City through a process of observation and field measurements and laboratory testing to obtain predetermined data parameters which are then used as a basis for pregaring river and groundwater quality status reports body in the Supiori district of Papua Province. Rusweri River is one of the largest rivers in Supiori Regency which has a river width of 90 m. Although this river is not a river intended for drinking water in Rusweri village, it is essential for the continuity of coastal habitats and ecosystems because the Rusweri river empties into the ocean. Upstream activities according to local community information contain local mines and do not use hazardous chemicals as an aid in obtaining gold, but the potential pressure of water resources can lead to inadequate water supply, deteriorating water quality, and low surface water flow<sup>3</sup>. The physical condition contains high dissolved solids originating from the river headwaters so that the water and river water quality monitoring activities are carried out on the Rusweri river in Supiori Regency, Papua Province.

Rasayan J. Chem., 14(1), 454-459(2021) http://dx.doi.org/10.31788/ RJC.2021.1416111



#### EXPERIMENTAL

The research site 33 he headwaters of the river up to the downstream of the Rusweri river, Supiori district, Papua Province. This research approach uses a qualitative approach. The method used in this study is a combination of quantitative. Quantitative methods, among others, compare the results of measurement as ameters in situ and laboratory and compare the data with water quality standards. The quantitative data will be calculated using the Pollution Index method. Another quantitative method used in determining the status of water quality using the Pollution Index method. Considerations using the Pollution Index method because there as no differences between the types of physical, chemical and biological contaminants. Pollution Index based on the Decree of the State Minister for the Environment No. 115 of 2003 concerning Guidelines for Determination of Water Quality Status. The Pollution Index for the allotment (j), which is a function of Ci / Lij can be determined by equation (1).

$$x = \frac{\sqrt{\left(\frac{Cij}{Lij}\right)}M^2 + \left(\frac{Ci}{Lij}\right)R^2}{2} \tag{1}$$

Where, Ci = Concentration of water quality parameters (i); Lij = Concentration of water quality parameters on water designation standards (j); Plj = Pollution Index / Pollution Index for designation (j); (Ci /  $\frac{1}{13}$ ) M = (Ci / Lij) maximum; (Ci / Lij) R = (Ci / Lij) average; Evaluate the value of PI (Pollution Index);  $0 \le PIj \le 1.0$  Meet quality standards (good condition);  $1.0 \le PIj \le 5.0$  Light pollution;  $5.0 \le PIj \le 10$  Medium polluted; PI> 10 Severe pollutants.

The Rusweri river water sampling location consists of 4 (four) locations along the river body. The location code and sample coordinates are as follows:

No	Location Codes	Codes and Sampling Coordinates  Coordinates		
1	S1	0040'26.94"S 135032'09.76"E		
2	S2	0040'32.03"S 135032'12.21"E		
3	S3	0040'36.62"S 135032'09.97"E		
4	S4	0040'40.91"S 135032'07.79"E		

### RESULTS AND DISCUSSION

Measurement of Rusweri river samples carried out in the laboratory is shown in Figs-2 to 10. The analysis conducted obtained results that refer to quality standards based on PP No. 82 of 2001. Discussion of parameters of river water quality parameters that exceed water quality standards and are shown in Figs.-1 to 7.




Fig.-1: Test Results for Phosphate Levels in the Rusweri River

Fig.-2: Test Content of Iron Content (Fe) in the Rusweri River




Fig.-3: Fluoride Test Results in the Rusweri River

Fig.-4: Sulphate (SO<sub>4</sub><sup>2-</sup>) Test Results in the Rusweri River




Fig.-5: TDS Test Results in the Rusweri River

Fig.-6: BOD Test Results in the Rusweri River




Fig.-7: COD Test Results in the Rusweri River

The test results show that of all parameters measured based on PP No. 82 of 2001, there were several chemical parameters that exceeded the established water quality standard, including Total Phospate location

#### RASĀYAN *J. Chem.* Vol. 14 | No. 1 |454-459| January - March | 2021

A1 =  $\frac{0}{.}655$  mg/L, A2 mg/L = 0.766 mg/L, A3 =  $\frac{0}{.}767$  mg/L, A4 = 1 mg/L, A5 =  $\frac{1}{100}$ g/L while PO<sub>4</sub> =  $\frac{0}{.}200$ mg/L quality standard; iron content (Fe) at location A3 = 0.3747 mg/L, A4 = 0.412 mg/L,  $\frac{1}{10}$  = 0.412 mg/L while the quality standard is 0.3 mg/L; Fluoride at location A4 = 0.8557 mg/L, A5 =  $12\frac{1}{28}$ /L while the standard quality is 0.5 mg/L; Sulfate levels (SO4) at location A4 = 487 mg/L and A5 = 547 mg/L the quality standard is 400 m 55; COD levels at locations A1 = 11 mg/L, A2 = 18.7 mg/L, A3 = 19.7 mg/L, A4 = 33.3 mg/L, A5 = 46.9 mg/L the quality standard is 0,2 mg/L; TDS levels for location A4 = 9.070 mg/L and A5 = 9.070 mg/L while the standard quality is 1,000 mg/L. Sulfur or sulfur is a chemical element with an atomic number of 16 represented by sulfur. In general, most of the sulfur ontained in the water is in the form of sulfate ions (SO<sub>4</sub><sup>2-</sup>). Sulfur (S) is in organic and inorganic form. Inorganic sulfur is mainly present in the form of sulfate (SO<sub>4</sub><sup>2-7</sup>), which is the main form of sulfur in waters and soils<sup>5</sup>. Sulfates that bind with hydrogen form sulfuric acid, and sulfates that bind to alkali metals are the most common form of sulfur found in lakes and rivers<sup>6</sup> Chemically sulfate is an inorganic form of sulfide in an aerobic environment. Scientifically, sulfate is derived from dissolving sulfur-containing minerals, for example, CaSO<sub>4</sub>·2H<sub>2</sub>O or casts and calcium sulfate anhydrous (CaSO<sub>4</sub>). WHO recommends that sulfate levels be permitted in drinking was around 400 mg/L and hydrogen sulfide levels around 0.05 mg/L<sup>7</sup>. The sulfate ion is a type of solidin with the empirical formula SO<sub>4</sub><sup>2</sup> with a molecular mass of 96.06 atomic mass units. Sulfates consist of a central sulfur atom surrounded by four oxygen atoms in a negatively charged two-sulfur tetrahedron arrangement8. The maximum limit of sulfate in water is around 250 mg/L for water consumed by humans<sup>9</sup>. The addition of Sulfate ions from brackish water makes it possible to increase the number of sulfate parameters at locations A3 and A4. The flow velocity of water around 0.2 m/s strongly supports the entry of a certain amount of water from the sea during high tide conditions increasing  $SO_4^{2-}$  levels at points A3 and A4. COD and BOD concentrations from upstream to downstream exceed the river water quality st 22 ard, and this data is strongly supported by the amount of oxygen dissolved in Rusweri river water (A1 = 6 mg/L, A2 = 5.3 mg/L, A3 = 4.3 mg/L, A4 = 4 mg/L and A5 = 3 mg/L) while the minimum standard quality is 6 mg / L. High phosphate levels in rivers can cause high levels in the sea if the river empties into the sea<sup>10</sup>, the source of phosphate in marine waters in coastal areas is rivers. The high levels of COD and BOD in the Rusweri river, when viewed from upstream to downstream, are caused by the decomposition of large amounts of organic matter in the waters which will absorb oxygen in the water thereby reducing the amount of dissolved oxygen (DO)11. The relationship between decreased DO levels with BOD and COD is shown in Figure 8.




Fig. -8: Relationship between Parameters [BOD] mg/L River Water, [COD] mg/L soil and [DO] mg/L

Iron (Fe) content in water at location A3 $^{16}$ 0.3747 mg/L, A4 = 0.412 mg/L, A5 = 0.412 mg/L while the standard quality of Fe in river water is 0.3 mg/L. The chemical condition of the soil around the river is a factor that affects Fe levels in water and the number of dissolved solids (TDS) in water<sup>12</sup> as illustrated in Fig.-9.




Fig.-9: Relationship between Parameters [Fe] mg/L River Water, [Fe] mg/Kg Soil and [TDS] mg/L

Rusweri river water quality status after being processed using the formula PIj (Pollution Index). Based on the calculation of the pollution index is shown in Table-2.

Table-2: Pollution Index Table based on Parameters in the Rusweri River Body

Parameter	Pollution Index	Status
PO <sub>4</sub>	5.417	Medium polluted
Fe	1.191	Mild polluted
F <sup>-</sup>	17.381	Heavily polluted
SO <sub>4</sub> 2 <sup>-</sup>	1.03	Mild polluted
TDS	6.925	Medium polluted
BOD	2.027	Mild polluted
		According to quality
COD	0.378	standards

Rusweri river water pollution index based on the calculation of PIj (pollution index) shown in Table-2, then the mild pollution status is more dominant, as shown in Fig.-10.




Fig.-10: Pollution Index based on number of PI status

#### CONCLUSION

This study concludes that the Rusweri river in Supiori Regency, Papua Province, is still consideral suitable for the development of ecosystems that exist in and around the river, the local community be personal some important parameters have exceeded the water quality standard. Urgently Rusweri river water can be consumed as raw water for drinking water but needs further processing.

The author would like to thank the Department of Chemistry, Gadjah Mada University, and Indonesia Endowment Fund for Education (LPDP-BUDIDN), for providing funds and facilities for this research.

#### REFERENCES

- 1. A.H. Mahvi, J. Nouri, A.A. Babaei, R. Nabizadeh, International Journal of Environmental Science and Technology, 2, 41(2005), DOI:10.3389/fpubh.2019.00174
- Y. Teng, J. Yang, R. Zuo, J. Wang, Journal of Earth Science, 22, 658 (2011), DOI:10.1007/s12583-011-0217-2
- 3. I.S. Saimy, N.A.M. Yusof, Procedia Social and Behavioral Sciences 81, 371(2013), **DOI:** 10.1016/j.sbspro.2013.06.445
- 4. D. Janardhana Rao, A.V.V.S. Swami, K. Raghu Babu, B. Hari Babu, Rasayan Journal of Chemistry, 9(3), 444 (2016).
- 5. Y. Chen, L. Shen, T. Huang, Z. Chu, Z. Xie, Science of the Total Environment, 703, 135591(2020), **DOI:** 10.1016/j.scitotenv.2019.135591
- 6. J.L. Duan, J.W. Sun, M.M. Ji, Y. Ma, Z.T. Cui, R.K. Tian, P.C. Xu, W.L. Sun, X.Z. Yuan, Microbiological Research, 236, 126453(2020), DOI:10.1016/j.micres.2020.126453
- 7. W.A.M. Fernando, I.M.S.K. Ilankoon, T.H. Syed, Minerals Engineering 117, 74(2018), DOI:10.1016/j.mineng.2017.12.004
- A.S. Thangiah, Rasayan Journal of Chemistry, 12, 1503(2019), DOI:10.31788/RJC.2019.1235201
- Y. Meride, B. Ayenew, Environmental Systems Research, 5(1), 1 (2016), DOI:10.1186/s40068-016-0053-6
- 10. A. Santhy, R. Rejithamol, Rasayan Journal of Chemistry, **11(3)**, 1067(2018), DOI:10.31788/RJC.2018.1131978
- 11. Y.M. Yustiani, S. Wahyuni, M.R. Alfian, Rasayan Journal of Chemistry, 11(2), 475(2018), DOI:10.7324/RJC.2018.1121892
- 12. D. Kumar, V. Kumar, S. Kumari, Rasayan Journal of Chemistry 11(3), 1477(2018), DOI:10.31788/RJC.2018.1143075

[RJC-6111/2020]

Masayan 2021	the influence of	3011	
ORIGINALITY REPORT			
27% SIMILARITY INDEX	19% INTERNET SOURCES	22% PUBLICATIONS	13% STUDENT PAPERS
PRIMARY SOURCES			
Submitte Student Paper	ed to United Int	ernational Col	llege 1 %
jurnal.ur	ntidar.ac.id		1 %
"Fisherie water qu district",	R M Milad, Y A es community b uality of the Dek IOP Conference mental Science,	ehaviour towa ket river, Lamo e Series: Earth	ards ongan
Ratnawa DEVELO MEMBRA	vi, Ghufira, S. Sa eti, D. A. Triawar PMENT OF PRO ANES BASED ON Journal of Cher	n, I. Gustian. "TON CONDUC N USED MATE	THE CTING
	, M. Padli, Tugiy NATION OF GAN		1 9

COMBINATION OF GAMBIER EXTRACT AND BENZOIC ACID AS INHIBITOR OF CALCIUM SULFATE SCALE FORMATION", Rasayan Journal of Chemistry, 2021

6	innspub.net Internet Source	1 %
7	Submitted to October University for Modern Sciences and Arts (MSA) Student Paper	1 %
8	essence-journal.com Internet Source	1 %
9	jestec.taylors.edu.my Internet Source	1 %
10	Marwan Khalish, Ayu Utami, Herwin Lukito, Susila Herlambang. "Evaluation of Textile Industry Wastewater Treatment as an Effort to Control River Water Pollution in Central Java", KnE Life Sciences, 2022	1 %
11	Submitted to Rutgers University, New Brunswick Student Paper	1 %
12	mdpi-res.com Internet Source	1 %
13	www.aes.bioflux.com.ro Internet Source	1 %
14	R M D Ujianti, S Anggoro, A N Bambang, F Purwanti. "Water quality of the Garang River, Semarang, Central Java, Indonesia based on	1 %

## the government regulation standard", Journal of Physics: Conference Series, 2018 Publication


15	academic-accelerator.com Internet Source	1%
16	businessdocbox.com Internet Source	1%
17	D. Justus Reymond, Karuppasamy Sudalaimuthu. "WATER QUALITY DURING PRE- MONSOON AND POSTMONSOON AND MODELLING OF TOTAL DISSOLVED SOLIDS FOR TAMIRAPARANI RIVER, TAMILNADU, INDIA", Rasayan Journal of chemistry, 2021 Publication	1 %
18	repo.jayabaya.ac.id Internet Source	1%
19	Ika Apriyani, Louise Elizabeth Radjawane. "Simulation Of Bod And Cod Pollutants Distribution In Tello River Makassar City", Journal of Physics: Conference Series, 2020 Publication	1%
20	Submitted to University of Central Oklahoma Student Paper	1 %
21	Submitted to Sriwijaya University Student Paper	1%

A. Sosa, M. Aurora Armienta, A. Aguayo, O. **1** % 22 Cruz. "Evaluation of the influence of main groundwater ions on arsenic removal by limestones through column experiments", Science of The Total Environment, 2020 Publication Amrita Das. "IMPACT OF SOME NATURAL 1 % 23 HAZARDS ON THE QUALITY OF WATER: RIVER DIANA, DISTRICT JALPAIGURI, WEST BENGAL, INDIA", Rasayan Journal of Chemistry, 2021 Publication Rohit Raj, K. S. Chandrashekar, Rounak 1 % 24 Biswas, Aravind Pai, Vasudev Pai. "RAPID HIGH-PERFORMANCE THIN LAYER CHROMATOGRAPHIC QUANTITATIVE ESTIMATION OF CAFFEINE IN VARIOUS FOODS AND BEVERAGES", Rasayan Journal of Chemistry, 2021 Publication Mustakim, Rosdina, Dian Ramadhani, M. <1% 25 Afdal, Medyantiwi Rahmawita. "The Classification Status of River Water Quality in Riau Province Using Modified K-Nearest

Riau Province Using Modified K-Nearest
Neighbor Algorithm with STORET Modeling
and Water Pollution Index", Journal of Physics:
Conference Series, 2021

Publication

and Environmental Science, 2021



Ade Lenty Hoya, Yuliastuti Nany, Sudarno Utomo, Andrea Ramadhani Maharlika. "Water quality assessment and control efforts use the pollution index method in the Sikendil river", E3S Web of Conferences, 2020

< 1 %

Auldry Walukow, I Nyoman Sukarta. "Analysis of Carrying Capacity and Water Pollution in the Simporo Strait Area After the Flash Flood", Ecological Engineering & Environmental Technology, 2021

< 1 %

Publication

Exclude quotes On Exclude bibliography On

Exclude matches

< 5 words