



DAVID PUBLISHING

From Knowledge to Wisdom


ISSN 2161-6256

DOI:10.17265/2161-6256

Journal of Agricultural Science and Technology A

Volume 9, Number 4, April 2019

Agricultural Resource; Plant Protection; Zootechny and Veterinary



David Publishing Company
www.davidpublisher.com

Journal of Agricultural Science and Technology A

Volume 9, Number 4, April 2019 (Serial Number 76)



David Publishing Company
www.davidpublisher.com

Publication Information:

Journal of Agricultural Science and Technology A (Earlier title: *Journal of Agricultural Science and Technology*, ISSN 1939-1250) is published monthly in hard copy (ISSN 2161-6256) by David Publishing Company located at 3 Germay Dr., Unit 4 #4651, Wilmington DE 19804, USA.

Aims and Scope:

Journal of Agricultural Science and Technology A, a monthly professional academic journal, particularly emphasizes new research results in agricultural resource, plant protection, zootechny and veterinary, all aspects of animal physiology, modeling of animal systems, agriculture engineering and so on. Articles interpreting practical application of up-to-date technology are also welcome.

Editorial Board Members (in alphabetical order):

| | | |
|---------------------------------|--|-------------------------------|
| Catherine W. Gitau (Australia) | Dharmatileke Bandula Kelaniyangoda (Sri Lanka) | |
| Ekachai Chukeatirote (Thailand) | Erin K. Espeland (USA) | Farzana Perveen (Pakistan) |
| Francesco Contò (Italy) | Francesco Montemurro (Italy) | Genhua Niu (USA) |
| Gulshan Mahajan (India) | Idress Hamad Attitalla (Libya) | Jang Ho Son (Korea) |
| Jagadish Timsina (Bangladesh) | Jelena Bošković (Serbia) | Manoj K. Shukla (USA) |
| Mehmet Musa Özcan (Turkey) | M. S. Qureshi (Pakistan) | Milad Manafi (Iran) |
| Mehmet Rüşü Karaman (Turkey) | Noureddine Benkeblia (Algeria) | Natraj Krishnan (USA) |
| Olivier A. E. Sparagano (UK) | Ozlem TOKUSOGLU (Turkey) | Renato S. Pacaldo (USA) |
| Ram C. Bhujel (Thailand) | Shoil M. Greenberg (USA) | Sanjeev Kumar Chauhan (India) |
| Shri Mohan Jain (Finland) | Thai Ngoc Chien (Vietnam) | T. Chatzistathis (Greece) |
| Vasudeo P. Zambare (USA) | Vasileios Fotopoulos (Greece) | Young Jung Kim (Korea) |
| Yusuf Bozkurt (Turkey) | Zeki Candan (Turkey) | |

Manuscripts and correspondence are invited for publication. You can submit your papers via web submission, or E-mail to agriculture@davidpublishing.org. Submission guidelines and web submission system are available at <http://www.davidpublisher.com>.

Editorial Office:

3 Germay Dr., Unit 4 #4651, Wilmington DE 19804, USA

Tel: 1-323-984-7526, 323-410-1082

Fax: 1-323-984-7374, 323-908-0457

E-mail: agriculture@davidpublishing.org, agriculture66@hotmail.com, agriculture@davidpublishing.com

Copyright©2019 by David Publishing Company and individual contributors. All rights reserved. David Publishing Company holds the exclusive copyright of all the contents of this journal. In accordance with the international convention, no part of this journal may be reproduced or transmitted by any media or publishing organs (including various websites) without the written permission of the copyright holder. Otherwise, any conduct would be considered as the violation of the copyright. The contents of this journal are available for any citation. However, all the citations should be clearly indicated with the title of this journal, serial number and the name of the author.

Abstracted/Indexed in:

Chemical Abstracts Service (CAS), USA

Ulrich's Periodicals Directory, USA

Chinese Database of CEPS, American Federal Computer Library Center (OCLC), USA

China National Knowledge Infrastructure (CNKI), China

Chinese Scientific Journals Database, VIP Corporation, Chongqing, China

Norwegian Social Science Data Services (NSD), Norway

Index Copernicus, Poland

Universe Digital Library Sdn Bhd (UDLSB), Malaysia

Google Scholar

PBN (Polish Scholarly Bibliography)

Electronic Journals Library

AcademicKeys

CiteFactor (USA)

Subscription Information:

Price (per year)

Print \$1200

David Publishing Company

3 Germay Dr., Unit 4 #4651, Wilmington DE 19804, USA

Tel: 1-323-984-7526, 323-410-1082 Fax: 1-323-984-7374, 323-908-0457

E-mail: order@davidpublishing.com

Digital Cooperative Company: www.bookan.com.cn



David Publishing Company

www.davidpublisher.com

Journal of Agricultural Science and Technology A

Volume 9, Number 4, April 2019 (Serial Number 76)

Contents

Review

- 201 **Synergistic Effects of Fumonisin B1 and B2 in Pigs: A Review**

Anilce de Araújo Brêtas and Patrícia Castelo Branco do Vale

Research Papers

- 207 **Monitoring Landslide Conditions in Madeira Island Using NOAA Operational Satellites**

Natalia Camara and Felix Kogan

- 227 **Controlling *Tuta absoluta* (Meyrick) by Selected Crude Plant Extracts in the Laboratory and in the Screen House**

Matendo Rehema Esther, Nonga Hezron Emmanuel, Bakari Gaymary George, Ndusha Nabintu Bintu, Mwatawala Maulid Walad and Maerere Amon Petro

- 240 **Feeding Effect of Different Levels of Agricultural and Food Waste on Growth Performance and Economics in Pig Production**

Trisiwi Wahyu Widayati, Bernadeta Wahyu Irianti Rahayu, Dwi Djoko Rahardjo and Marlyn Nelce Lekitoo

- 248 **Preliminary Researches Regarding the Effectiveness of the Formic Acid Treatment on *Varroa* (*Varroa destructor*) Found in the Artificially Decapped Bee Brood**

Adrian Siceanu, Eliza Căuia, Gabriela Oana Vișan and Dumitru Căuia

- 262 **Diversity of Stingless Bees Based on Principal Component Analysis in Halmahera Island, Indonesia**

Hearty Salatnaya, Asnath Maria Fuah, Cece Sumantri, Widiatmaka and Sih Kahoho

Feeding Effect of Different Levels of Agricultural and Food Waste on Growth Performance and Economics in Pig Production

Trisiwi Wahyu Widayati, Bernadeta Wahyuni Irianti Rahayu, Dwi Djoko Rahardjo and Marlyn Nelce Lekitoo
Department of Animal Science, University of Papua, Jl. Gn. Salju Amban Manokwari Papua Barat 98314, Indonesia

Abstract: This study was undertaken with the objectives to determine the feeding effect of different levels of agricultural and food waste on growth performance and cost effectiveness in pig production. This study was conducted at Manokwari regency, West Papua Province, Indonesia. The agricultural and food industry by-products used were collected from two traditional markets, five restaurants and 15 small-scale food industries. All materials used as ration were proximally analyzed to determine their nutritional values. Average daily gain (ADG), feed consumption and feed conversion ratio (FCR) were taken to determine the pigs' performance. Feed cost using local market prices was estimated in knowing the ability of agricultural waste product and food industry to substitute commercial feed. Tabulation was used to analyse the data. The result of this research indicates that the usage of agricultural and food industries' wastes as the alternative components of pigs' feed to substitute the commercial ones does not reduce the pigs' performance and production. Among three rations that use agricultural and food industry wastes, ration with the combination of 25% waste and 75% commercial feed had the best FCR. On the other hand, a mixture of 75% waste and 25% commercial feed is the most economic one, and able to reduce the cost of feeding up to 35.86%.

Key words: Agricultural by-product, pig, ration, ration cost.

1. Introduction

The paradigm of modern animal husbandry is based on zero waste concept. Use of agricultural and food industry by-products into useful materials is an important issue that needs to be done.

In the last 10 years many researchers have paid attention to the existence of waste both from agriculture and by-product from the food industry. Without proper management, wastes both from agricultural and food industry are considered to be a threat both ecologically and aesthetically to the environment and human health. Obi *et al.* [1] revealed that agricultural wastes are defined as the residues from the growing and processing of raw agricultural product as fruits, vegetable, meat, poultry dairy product and crop. Meanwhile, industry by-products are produced in

the large amount in the food industry. Helkar *et al.* [2] state that about 38% of food wastes occur during food processing. Food wastes are produced by a variety of sources, animal-derived processing food wastes include by-products from breed animals such as carcasses, hides, hoofs, heads, feathers, manure, offal, viscera, bones, fat and meat trimmings, blood; wastes from seafood such as skins, bones, oils, blood; wastes from dairy processing industry such as whey, curd, and milk sludge from the separation process; vegetable-derived processing food wastes include peelings, stems, seeds, shells, bran, trimmings residues after extraction of oil, starch, juice and sugars. Improper management of foodstuffs also reduces the total economic value of food. Venkat [3] stated that the loss of economic value of food such as vegetables, fish, legumes caused by inefficiency in retail and consumer in America was as much as US\$197.68 billion per year. Therefore, waste management is

Corresponding author: Trisiwi Wahyu Widayati, Ph.D., research field: social and economic of agriculture.

needed to increase the value of the waste. In some countries with a high demand for animal feeding such as Japan, South Korea and Taiwan, local law encourages using food waste to feed animal [4].

Papua is one of the provinces in the eastern part of Indonesia with the indigenous population called Papuan. Papuan has a unique culture and social life where they make pigs as the favorite animal. Pigs for the people of Papua are valuable in social, cultural and economic setting. Most of the communities use pigs in traditional ceremonies, and also as an exchange tool in several transactions. The market demand of this commodity is quite high and it has become a primary saving for many farmers. The selling price of this animal is sufficiently high, ones in weaning period can be sold from IDR 1,000,000 to IDR 1,500,000 (\$70.88 to \$106.32) and the price for ones aged 8-12 months varied from IDR 3,000,000 to IDR 5,000,000 (\$212.64 to \$354.40). Unfortunately so far, pigs in Papua have not been intensively raised yet. The animals usually are left to look for food by themselves, so the aspects of feeding, reproduction and health are not properly cared. In general, farmers feed their animal with only a single type of feed, such as tubers or roots which are low in quality. The minimum amount and the low quality of feed are factors that affect the slow growth of pig and make them more vulnerable to diseases [5, 6]. Another problem in raising pigs is that they are a monogastric animal and many of their feed competed with human's food. This condition causes a problem in feed availability during intensive pig raising, thus it is imperative to find alternatives to high quality pigs' feed without competing with the people's [7, 8]. According to Ref. [9], West Papua has an ample amount of forages and agricultural by-products and has the potential to sustain the development of animal husbandry, which is 42,442,750 t produced from 4,244,274 ha. Furthermore, Pattiselanno and Iyai [10] reported that food wastes from restaurants, food industries and traditional fish markets were 2,056, 200 and 500 kg/d, respectively.

A good livestock development is dependent upon the availability of feed, socio-cultural conditions and local climate [11, 12]. In fact, the potential market of pig is high, but on the other hand there are still problems in the continuous feed: whether it is availability, quality or cost. Thus it is necessary to use the ingredients from agriculture and food industry by-products as an alternative of pig ration [13, 14]. Based on reasons above, a study was conducted to evaluate potential of agriculture and food industry by-products in Manokwari regency as nutrient source to pig and its ability to substitute commercial ration in order to reduce ration cost.

2. Materials and Methods

2.1 Location of Study

The experiment was carried out from June to August 2017 at the Animal Research Laboratory of University Papua in Manokwari, Indonesia (134°04' longitude and 00°48' latitude) with a mean altitude of 110 m. Average temperature is 37 °C with daily variations from 37-39 °C.

2.2 Animal, Ration and Experimental Design

Twelve (12) male local pigs with an initial body weight (BW) of 12.54 ± 1.84 kg were arranged in a completely randomized design (CRD) with four treatments and three replications. The animals were housed in 12 individual cages.

Ration used in this experiment were formulated from agricultural and food industry by-products, i.e., fish waste, soybean curd waste, taro skin, soybean skin, vegetables waste and commercial pig ration. The agricultural and food industry by-products were collected from two traditional markets, five restaurants and 15 food industries in Manokwari regency. The four treatments were T1: combination of 75% agricultural and food industry by-products and 25% commercial ration; T2: combination of 50% agricultural and food industry by-products and 50% commercial ration; T3: combination of 25% agricultural and food industry

by-products and 75% commercial ration; T4: 100% commercial ration. Rations T1, T2 and T3 were formulated based on dry matter (DM) basis with iso-protein and iso-energy concept. The commercial pig ration (Charoen Pokphand 511) as control diet used in this experiment was produced by Charoen Pokphand, Indonesia. Formulation of pig ration in starter period is presented in Table 1.

2.3 Experimental Procedure

The experiment lasted 35 d and was comprised of 10 d for adaptation period, followed by four weeks for feed intake data collection. The rations supplied twice a day (at 08:00 and 16:00 h) *ad libitum*. Fresh water is available *ad libitum*. Individual ration refusals, if any, were collected, weighed daily and samples were collected for analysis. Before the start of the experiment, pigs were dewormed with 0.7 mL/kg BW of albendazole (PT. Kimia Farma, Indonesia). The animals were weighed each week throughout the experiment.

2.4 Statistical Analysis

Data were subjected to analysis of variance for a CRD using SPSS version 21. Comparison of means was carried out using the Duncan's multiple range tests, when the effect of treatment was significant ($p < 0.05$).

The linear model of experimental design was as follows:

$$Y_{ij} = \mu + T_i + \varepsilon_{ij}$$

where Y_{ij} : the score for observation of i th variable and

j th replication; μ : the overall population mean; T_i : the effect of i th treatment level (type of ration); ε_{ij} : the error effect associated with i th treatment level and j th replication; i : 1, 2, 3, 4; j : 1, 2, 3.

3. Results

The agricultural and food industry by-product availability in Manokwari regency is shown in Table 2. The utilization of agricultural and food industry wastes is expected to improve their economic value as pigs' feed.

The most important factor in livestock's ration formulation is the balance between energy and protein composition. That is why proximate analysis was conducted, with the result displayed in Table 3.

In the farm business, the feeding part is the most expensive of all the operational cost. A study conducted by Sala and Delia [15] indicated that the feeding part covered 80% of the livestock upkeep cost. As shown in Table 4, waste utilization was able to reduce the feeding cost, the most cost reduction was achieved with 75% of waste mixture.

Three kinds of treatments were made from wastes, while the last one was commercial feed as the control. Tabulation and statistic tests results of agricultural waste usage as starter pigs' feed are presented in Table 5.

The result indicated that the utilization of all four kinds of ration was able to increase the pigs' daily weight gain, with the highest average from T3 ration (0.465 ± 0.05 kg daily) followed by T4, T2 then T1.

The analysis of variance showed that there were no

Table 1 The composition of pig rations (%) in starter period.

| Feedstuffs | T1 | T2 | T3 | T4 |
|--------------------|-------|-------|------|-----|
| Fish waste | 18.57 | 12.37 | 6.17 | |
| Soybean curd waste | 15.70 | 10.47 | 5.23 | |
| Taro skin | 8.82 | 5.9 | 2.94 | |
| Vegetables waste | 16.04 | 10.69 | 5.37 | |
| Restaurant waste | 15.87 | 10.57 | 5.29 | |
| Commercial ration* | 25 | 50 | 75 | 100 |
| Total | 100 | 100 | 100 | 100 |

* Contained corn, rice bran, soybean meal, coconut meal, meat and bone meal, wheat meal, canola, calcium, phosphorus, vitamin, trace mineral and anti-oxidant.

Table 2 Types and potentials of local feed in Manokwari regency.

| No. | Feed commodity | Potential (kg/year) ^a | By-products | Potential (kg/d) ^b |
|-----|----------------|----------------------------------|------------------------------------|-------------------------------|
| 1 | Soybean | 15,191.00 | Soybean curd waste Soybean skin | 2,400.00 55.50 |
| 2 | Fish | 27,911.00 | Fish waste | 1,000.00 |
| 3 | Mung bean | 1,891.00 | Mung bean skin | 83.40 |
| 4 | Rice | 115,865.80 | Rice bran | 11,586.60 |
| 5 | Bananas | 9,441.10 | Bananas skin | 127.50 |
| 6 | Taro | | Taro skin | 11.40 |
| 7 | Vegetables | 4,345.20 | Vegetables waste | 546.00 |
| 8 | Restaurant | 1,521 | Restaurant waste | 2,056.06 |

^a BPS-Statistic of Papua Barat Province (2014) [16]; ^b Primary data of survey.

Table 3 The potential and nutrients content of ingredients in pig ration.

| No. | Ingredients | Nutrients content | | | |
|-----|--|---------------------|-------------------------------------|--|--|
| | | Dry matter (DM) (%) | Crude protein (CP) (%) ^a | Gross energy (GE) (kcal/kg) ^a | Metabolizable energy (ME) (kcal/kg) ^b |
| 1 | Fish waste | 29.41 | 31.21 | 3,432.94 | 2,709 |
| 2 | Soybean curd | 14.31 | 23.85 | 4,950.57 | 3,906 |
| 3 | Soybean skin | 15.96 | 15.1 | 4,022.23 | 3,174 |
| 4 | Taro skin | 26.45 | 4.26 | 3,648.96 | 2,879 |
| 5 | Vegetables waste | 9.84 | 15.8 | 3,683.99 | 2,907 |
| 6 | Restaurant waste | 35.84 | 13.72 | 4,202 | 3,315 |
| 7 | Commercial pig ration (Charoen Pokphand 511) | 87 | 19.5 | - | 3,315.12 |

^a DM basis; ^b Based on calculation.

Table 4 Feeding cost reduction of each treatment compared to commercial ration.

| No. | Feedstuffs | Prices (IDR/kg) | T1 | | T2 | | T3 | | T4 | |
|-----|--|-----------------|----------------------|------------|----------------------|------------|----------------------|------------|----------------------|------------|
| | | | As fed (kg/head/day) | Cost (IDR) | As fed (kg/head/day) | Cost (IDR) | As fed (kg/head/day) | Cost (IDR) | As fed (kg/head/day) | Cost (IDR) |
| 1 | Fish waste | 1,250 | 0.196 | 245 | 0.124 | 155 | 0.064 | 80 | | |
| 2 | Soybean curd | 800 | 1.074 | 859 | 0.683 | 546 | 0.354 | 283 | | |
| 3 | Taro skin | 1,400 | 0.309 | 433 | 0.197 | 276 | 0.102 | 143 | | |
| 4 | Vegetable waste | 1,500 | 1.613 | 2,420 | 1.025 | 1,538 | 0.534 | 801 | | |
| 5 | Restaurant waste | 500 | 0.426 | 213 | 0.27 | 135 | 0.14 | 70 | | |
| | Commercial ration Charoen Pokphand 551 | 11,000 | 0.263 | 2,897 | 0.502 | 5,524 | 0.782 | 8,599 | 1 | 11,000 |
| | Total feed | | 3.882 | 7,067 | 2.802 | 8,175 | 1.977 | 9,977 | 1 | 11,000 |
| | Cost reduction (%) | | | 35.76 | | 25.69 | | 9.3 | | 0 |

T1: combination of 75% agricultural and food industry by-products and 25% commercial ration; T2: combination of 50% agricultural and food industry by-products and 50% commercial ration; T3: combination of 25% agricultural and food industry by-products and 75% commercial ration; T4: 100% commercial ration.

Feeding Effect of Different Levels of Agricultural and Food Waste on Growth Performance and Economics in Pig Production

Table 5 Effect of agricultural and food industry by-products in ration on average daily gain (ADG), feed intake, feed conversion and feed cost.

| Variables | Treatments | | | | <i>p</i> -value | Sig. |
|-----------------------------|--------------------|--------------------|--------------------|---------------------|-----------------|------|
| | T1 | T2 | T3 | T4 | | |
| ADG (kg/head/day) | 0.444 ± 0.01 | 0.445 ± 0.03 | 0.465 ± 0.05 | 0.457 ± 0.02 | 0.791 | ns |
| Feed intake (kg/head/day) | 1.054 ± 0.01 | 1.004 ± 0.05 | 1.042 ± 0.04 | 1.002 ± 0.05 | 0.353 | ns |
| Feed conversion ratio (FCR) | 2.375 ± 0.06 | 2.259 ± 0.09 | 2.25 ± 0.13 | 2.193 ± 0.02 | 0.159 | ns |
| Feed cost (IDR/kg) | 7.066 ^a | 8.174 ^b | 9.976 ^c | 11.017 ^d | 0.000 | ** |

T1: combination of 75% agricultural and food industry by-products and 25% commercial ration; T2: combination of 50% agricultural and food industry by-products and 50% commercial ration; T3: combination of 25% agricultural and food industry by-products and 75% commercial ration; T4: 100% commercial ration.

Means with different superscripts in the same row differ significantly ($p < 0.01$); ** $p < 0.01$; ns: non-significant.

significant differences among treatments, whether it was T1, T2, or T3 against T4 (the control) for the pigs' performance, which were denoted by daily weight gain, feed intake, and feed conversion.

4. Discussion

The re-utilization of agricultural and food industry is expected to not only improve the environment but also to increase the wastes' economic value.

Wastes utilizations as shown in Table 1, were adjusted to crude protein (CP) and metabolizable energy (ME) requirements for starter-pigs' metabolism which were 19.5% and 3,150 kcal/kg. The feed used was vegetable waste, fish waste, tofu waste, taro waste and wastes from restaurants. The reason for the utilization of those wastes was that they were easily obtained in Manokwari. Two kinds of ingredients which were fish waste and soybean curd used in this study were included as protein sources, while others such as vegetable waste, restaurants' waste and taro skin were used as the energy sources. Taro skin had the lowest CP content (4.26%), otherwise the highest CP content obtained was from fish waste (31.21%). Tofu waste was the leftovers from the tofu-making process and weighed around 25%-35% of the final tofu product. The tofu waste used in this research had 23.85% CP content, this was consistent with the statement of Mathius and Sinurat [16] that tofu waste could be utilized as protein source feed since they had high gross protein value, around 23%-29%. The commercial feed (Charoen Pokphand 511) was used in this research as the control.

Growth is defined as the interaction between genetic, food and the environment [18]. Livestocks' growth as the result of efficient maintenance could be measured by feed intake, body-weight gain and the FCR [19, 20]. The body-weight gain of starter-phase pigs fed with four different feed is shown in Fig. 1.

As seen in Fig. 1, the pigs fed with T1 and T2 had more fluctuate graph compared to the ones fed with T3 and T4. The body-weight gains of T1 and T2 treatments were declining in the first two weeks, it was because the pigs were adapting to the new feed. Furthermore, during the period between the second and third weeks, the weight gain increased, but it went down again during the period between the third and fourth weeks. On the other hand, the weight gains for the T3 and T4 (control) treatments had a linear continuous growth from the first to the fourth weeks.

The weight fluctuation of the pigs with T1 and T2 treatments were thought to be caused by their monogastric nature, which lacked the ability to utilize feed with high amount of crude fiber content [20]. Because of that, even when the feed was made with the concept of iso-protein and iso-energy in mind, the lack of cellulose enzyme was the reason why not all of the feed was fully digested. However several studies have demonstrated that when pigs are feed with raw fiber high content of cellulose and lignin, there was a positive influence on their health status [14].

In feed intake aspect, the highest intake was seen in pigs fed with T1, followed by T2, T3 and lastly T4. Allegedly, it was because of the relatively higher palatability of agricultural and food industry wastes.




Fig. 1 The average body weight of pig during experiment.

Feed with many wastes was more liked by the pigs. This matched with a study conducted by Wea [22] about the usage of market wastes such as water spinach, fish waste and chaff as a mixture for pigs' ration, in which the more they are mixed, the higher the consumption rate.

Feed conversion is the amount of feed which needed to be consumed to gain 1 kg of BW. The lower the value, the higher the efficiency of the feed to increase the livestock's BW [23]. The lowest to the highest conversion rates were T4 (100% commercial feed), followed by T3, T2 and T1. This fits with studied Ref. [24] which stated that livestock feed in form of pellets was superior in terms of digestibility, because they need less energy to be digested. Therefore, to improve pigs' ability to digest feed made from agricultural and food industry wastes in the future, it is necessary to consider the production process, to determine which kind of size or form that is easier to digest.

The result of a research done by Warouw *et al.* [25] and Kueain *et al.* [26] showed that feeding cost took around 44.66% to 55% of all production cost. Such high percentage could be reduced by finding alternative source of feed, which will not reduce the output quality. The result of the research regarding

feeding cost is shown in Table 4, using three kinds of waste-based rations, proved that T1 ration could reduce the feeding cost up to 35.86%, T2 25.80% and T3 could reduce the cost compared to commercial feed by 9.45%.

In Manokwari, the commercial ration's cost is IDR 11,000 (\$0.77)/kg. However, the availability is not continuous yet. Considering the potential of market and food industry wastes are adequate, it is reasonable to strive to process those wastes to provide for the continuous availability of livestock feed. Cooperation with the local government is needed to build a collection center for the wastes from the markets' hub. The collection process will simplify further processing to create livestock feed which has economic value while bolstering the environment's hygiene.

5. Conclusions

The results of this study showed that there are no significant differences between the pigs' performance, whether they are fed with agricultural and food industry wastes or with commercial feed. Feeding treatment with the combination of 25% waste and 75% commercial feed gave the highest FCR average. On the other hand, the mixture of 75% waste and 25% commercial feed had the most economic price, and

could reduce the feeding cost up to 35.86%. Due to no significant difference between those treatments, the farmers should not hesitate to use the mixture of 75% waste and 25% commercial feed.

Acknowledgment

This study was funded by the Directorate General of Strengthening Research and Development, Indonesian Ministry of Research, Technology and Higher Education through “Hibah Produk Terapan” scheme (contract number 089/SP2H/LT/DRPM/IV/2017).

References

- [1] Obi, F. O., Ugwuishiwu, B. O., and Nwakaire, J. N. 2016. “Agricultural Waste Concept, Generation, Utilization and Management.” *Niger. J. Technol.* 35: 957-64.
- [2] Helkar, P. B., Sahoo, A. K., and Patil, N. J. 2016. “Review: Food Industry by Products Use as a Functional Food Ingredient.” *Int. J. Waste Resour.* 6: 248.
- [3] Venkat, K. 2011. “The Climate Change and Economic Impact of Food Waste in the United States.” *Int. J. Food Syst. Dyn.* 2: 431-46.
- [4] Thi, N. B. D., Kumar, G., and Lin, C.-Y. 2015. “An Overview of Food Waste Management in Developing Countries: Current Status and Future Perspective.” *J. Environ. Manage.* 157: 220-9.
- [5] Iyai, D. A., Rahayu, B. W. I., Sumpe, I., and Saragih, D. 2011. “Analysis of Pig Profile on Small-Scale Pig Farmer in Manokwari West Papua.” *J. Indon. Trop. Anim. Agric.* 36: 190-6.
- [6] Harikumar, S., Davis, J., and Anil, K. S. 2016. “Use of Unconventional Feed in Pig Production.” *J. Indian Vet. Assoc.* 14: 5-11.
- [7] Woyengo, T. A., Beltranena, E., and Zijlstra, R. T. 2014. “Controlling Feed Cost by Including Alternative Ingredients into Pig Diet: A Review.” *J. Anim. Sci.* 92: 1293-305.
- [8] Schader, C., Muller, A., Scialabba, N. E., Hect, J., Isensee, A., Erb, K. H., Smith, P., Makkar, H. P. S., Leiber, F., Schwegler, P., Stolze, M., and Niggli, U. 2015. “Impact of Feeding Less Food Competing Feedstuffs to Livestock on Global Food System Sustainability.” *J. R. Soc. Interface* 12: 1-12.
- [9] Statistic of Papua Barat Province (BPS). 2007. *Papua Barat Province in Figure*. BPS-Statistic of Papua Barat Province.
- [10] Pattiselanno, F., and Iyai, D. A. 2005. “Useful Waste: Pig Farmers’ Hope for Feed Source Alternatives in Manokwari’s Coastal Region.” *Salam* 13: 24-5.
- [11] Jabbar, M. A., and Akter, S. 2008. “Market and Other Factors Affecting Farm Specific Production Efficiency in Pig Production in Vietnam.” *J. Int. Food Agribus. Mark.* 20: 29-53.
- [12] Babovic, J., Caric, M., Djordjevic, D., and Lazic, S. 2011. “Factors Influencing the Economic of the Pork Meat Production.” *Agric. Econ.* 57: 203-9.
- [13] Kasapidou, E., Sossidou, E., and Mitlianga, P. 2015. “Fruit and Vegetable Co-products as Functional Feed Ingredients in Farm Animal Nutrition for Improved Product Quality.” *Agric.* 5: 1020-34.
- [14] Katsoulis, K., Leontide, L., and Kontopidis, G. 2016. “Locally Produced Agricultural by Product as Feed Sources for Pigs.” *J. Vet. Sci. Med.* 4: 1-5.
- [15] Sala, F., and Delia, E. 2012. “The Effect of Pellet Feed on the Performance and Nutrient’s Digestibility of Weaned Piglets.” *Albanian J. Agric. Sci.* 4: 275-8.
- [16] Statistic of Papua Barat Province (BPS). 2014. *Papua Barat Province in Figure*. BPS-Statistic of Papua Barat Province.
- [17] Mathius, I. W., and Sinurat, A. P. 2001. “Utilization of Unconventional Feedstuffs for Animal Production.” *Wartazoa* 11: 20-31.
- [18] Hardjosubroto, W. 1994. *Field Breeding Application*. Jakarta: Gramedia Widiasarana Indonesia.
- [19] Devi, S. M., Devi, U. S., and Kim, I. H. 2014. “Evaluation of Dietary Sources of Protein on Growth Performance in Pigs.” *Veterinari Med.* 59: 247-53.
- [20] Racadembosch, J., Amador, J., Bernaus, J., Font, J., and Fraile, L. J. 2016. “Production Parameter and Pig Production Cost; Temporal Evolution 2010-2014.” *Porcine Health Manage.* 2: 1-11.
- [21] Mace, O. J., and Marshall, F. 2013. “Digestive Physiology of the Pig Symposium: Gut Chemosensing and the Regulation of Nutrient Absorption and Energy Supply.” *J. Anim. Sci.* 91: 1932-45.
- [22] Wea, R. 2017. “Production Performance of Pigs Which Consume Market’s Organic Wastes.” *Partner J.* 20: 127-35.
- [23] Metzler-Zebeli, B. U., Lawlor, P. G., Magowan, E., McCormack, U. M., Curiaño, T., Hollmann, M., Ertl, R., Aschenbach, J. R., and Zebeli, Q. 2017. “Finishing Pigs That Are Divergent in Feed Efficiency Show Small Differences in Intestinal Functionality and Structure.” *PlosOne* 12: 1-16.
- [24] Massuquetto, A., Durau, J. F., Schramm, V. G., Netto, M. V. T., Krabbe, E. L., and Maiorka, A. 2018. “Influence of Feed Form and Conditioning Time on Pellet Quality, Performance and Ileal Nutrient Digestibility in Broilers.” *J. Appl. Poult. Res.* 27: 51-8
- [25] Warouw, Z. M., Panelewen, V. V. J., and Mirah, A. D. P.

2014. "Pig Farm Analysis of Kasewean Company at Kakaskasen II of Tomohon Municipality." *J. Zootek* 34: 92-102.
- [26] Kueain, Y. A., Suamba, I. K., and Wijayanti, P. U. 2017. "Business Financial Analysis on Piggery (A Case Study of Piggery of UD Karang at Jagapati Village, Sub-District of Abiansemal, Badung)." *Agribusiness and Agro-tourism Journal* 6: 96-104.



Journal of Agricultural Science and Technology A

Volume 9, Number 4, April 2019

David Publishing Company

3 Germay Dr., Unit 4 #4651, Wilmington DE 19804, USA

Tel: 1-323-984-7526, 323-410-1082; Fax: 1-323-984-7374, 323-908-0457

<http://www.davidpublisher.com>, www.davidpublisher.org

agriculture@davidpublishing.com, agriculture@davidpublishing.org

ISSN 2161-6256



9 772161 625194