

Data on environmental contaminants in sea turtle eggs at Venu Island, Kaimana – West Papua, Indonesia

by Rima Siburian

Submission date: 04-Oct-2020 04:42AM (UTC-0400)

Submission ID: 1404579518

File name: 1-s2.0-S2352340920306727-main.pdf (226.58K)


Word count: 1820

Character count: 10049



Data Article

Data on environmental contaminants in sea turtle eggs at Venu Island, Kaimana – West Papua, Indonesia



Ricardo F Tapilatu^{a,b,*}, Hengki Wona^a, Rima H. S Siburian^{a,c}

^a Research Centre of Pacific Marine Resources, University of Papua (UNIPA), Manokwari, 98314 Papua, Indonesia

^b Marine Science Department, University of Papua (UNIPA), Manokwari, 98314 Papua, Indonesia

^c Silviculture Department, University of Papua (UNIPA), Manokwari, 98314 Papua, Indonesia

ARTICLE INFO

Article history:

Received 28 April 2020

Revised 14 May 2020

Accepted 19 May 2020

Available online 27 May 2020

Keywords:

Contaminants

Sea turtle egg

Human consumption

Food chain

ABSTRACT

Long-range atmospheric transport of contaminants can result in contamination of aquatic systems in otherwise pristine areas. Contaminants enter aquatic systems largely as inorganic pollutants that often are microbially transformed. The result can be a potent neurotoxin with a strong tendency to bio-magnify within food chains. The general human population is exposed to contaminants such as mercury, predominantly through food consumption. Here, we report data on contaminants from a sample of green and hawksbill turtle that were sampled from a subset of two nests during the 2016 nesting season at Venu Island, Kaimana. Three contaminants (mercury, cadmium, arsenic) found in eggs exceeded safety limits for human consumption. Other contaminants such as lead, zinc, manganese, copper and iron were found below the established safety standard.

© 2020 Elsevier Inc.

This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

* Corresponding author at: Research Centre of Pacific Marine Resources, University of Papua (UNIPA), Manokwari, 98314 Papua, Indonesia.

E-mail address: rf.tapilatu@unipa.ac.id (R.F. Tapilatu).

<https://doi.org/10.1016/j.dib.2020.105778>

2352-3409/© 2020 Elsevier Inc. This is an open access article under the CC BY-NC-ND license.

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Specifications table

| | |
|--------------------------------|---|
| Subject | Pharmacology, Toxicology and Pharmaceutical Science: Toxicology |
| Specific subject area | Environmental Contaminants in Sea Turtle Eggs |
| Type of data | Data of contaminants in Table |
| How data were acquired | GBC 906 AA model atomic absorption spectrophotometer complete with inbuilt window-based 906 program software was used in this analysis |
| Data format | Raw [Kind and content of contaminants in sea turtle eggs] |
| Parameters for data collection | Contaminants (mercury, cadmium, arsenic, lead, zinc, iron, and copper) in sea turtle eggs |
| Description of data collection | Eggs were analysed for contaminants using an Atomic Absorption Spectrometer (AAS) |
| Data source location | Institution: Research Centre of Pacific Marine Resources (RCPMR) – University of Papua (UNIPA) City/Town/Region: Venu Island, Kaimana – South of Bird's Head Seascape – West Papua Country: Indonesia Latitude and longitude: 04° 7'35S; 134° 36'40E |
| Data accessibility | Repository name: University of Papua (UNIPA) Library Data identification number: 063.PS/X/2016 Direct URL to data: http://repository.unipa.ac.id:8080/xmlui/handle/123456789/433 |
| Related research article | Ricardo F. Tapilatu, Hengki Wona and Petrus P. Batubara. 2017. Status of sea turtle populations and its conservation at Bird's Head Seascape, Western Papua, Indonesia. <i>Biodiversitas</i> Vol. 18, Number 1, January 2017 Pages: 129-136. DOI: 10.13057/biodiv/d180119 |

Value of the data

- These data are the first reported environmental contaminants in sea turtle eggs in Kaimana, south of the Bird's Head Seascape (BHS) – Papua, Indonesia.
- These data are beneficial to provide recommendations that harvesting of turtle eggs and adults and juveniles of all four species of sea turtle at Kaimana be totally prohibited.
- These data have value. The health risk of consuming eggs contaminated with chemical compounds could potentially reduce the number of eggs collected for human consumption.
- These data allow further analysis to understand the connections between the foraging and nesting habitats and the pelagic phase of sea turtles in Kaimana and their relationships to contaminants.

1. Data description

Kaimana is one of several areas of the Bird's Head Seascape that sea turtles use during the nesting season [1]. Local villagers reported seeing sea turtles nesting on beaches in Lakahia and Venu Island. Sea turtles in Kaimana likely face a high risk of being harvested because of the nesting beach's proximity to human settlements [1]. In environmental assessments of heavy metals: cadmium and mercury warrant special attention because of their vast global distribution and high potential toxicity and carcinogenicity in humans. Data from numerous studies of these metals in sea turtles have indicated that their concentrations vary greatly by species, region, and tissue type. Cadmium, mercury, and lead have been documented in sea turtle eggs and hatchlings in concentrations known to cause toxic effects in other vertebrates [2,3].

The mercury content found in eggs from two subsets of nests exceeds safety limits for human consumption, 0.5 mg/kg [4] (Table 1). The World Health Organisation has adopted the United States Environmental Protection Agency guideline levels for mercury and recommends that food with mercury concentrations of 0.5 mg/kg or more should not be sold for human consumption. This suggests that the levels found in sea turtle eggs (up to 8.15 ppm) could be hazardous to human health. The cadmium levels found in eggs from five subsets of sea turtle nests from Venu Island exceeded established safety standards for seafood products, 0.5 mg/kg [4] (Table 1) and

Table 1

Assessment of contaminant content in a subset of sea turtle eggs collected from nests at Venu Island.

| No | Parameter | Detection Limit | Level detected in green turtle eggs (unit: mg/kg) | Level detected in hawksbill turtle egg (unit: mg/kg) | Analysis Method |
|----|----------------|-----------------|---|--|---------------------------|
| 1. | Mercury (Hg) | 0.004 | 0.421* | 0.326* | APHA, 2012 3112-B, 3030-E |
| 2. | Cadmium (Cd) | 0.40 | 2.12* | 2.31* | APHA, 2012 3111-B, 3030E |
| 3. | Arsenic (As) | 0.003 | 0.875* | 0.569* | APHA, 2012 3114-B, 3030E |
| 4. | Lead (Pb) | 0.23 | <0.23 | <0.23 | APHA, 2012 3111-B, 3030-E |
| 5. | Zinc (Zn) | 0.67 | 70.36 | 58.83 | as above |
| 6. | Manganese (Mn) | 0.53 | <0.53 | 0.80 | as above |
| 7. | Copper (Cu) | 1.2 | 4.33 | 3.65 | as above |
| 8. | Iron (Fe) | 5.00 | 15.65 | 42.78 | APHA, 2012 3500-Fe-B |

* Exceeding safety limits.

may warrant concern for people who consume sea turtle tissue with high cadmium concentrations in the Kaimana region. In addition, arsenic levels found in eggs from five subsets of sea turtle nests from Venu Island exceeded established safety standards for seafood products, 0.5 mg/kg [4] (Table 1) and may warrant concern for people who consume sea turtle tissue with high arsenic concentrations in the Kaimana region.

Other contaminants such as lead, zinc, manganese, copper and iron were found below the established safety standard (Table 1).

2. Experimental design, materials, and methods

Large scale contaminants studies of sea turtles eggs have often been avoided because they require the sacrifice of eggs in a threatened and/or endangered population [5]. In 2016, three fresh sea turtle eggs each from a recently laid nest of green (*Chelonia mydas*) and hawksbill (*Eretmochelys imbricata*) sea turtles were collected randomly to be analyzed for contaminants (Table 1). The six collected eggs were stored in a refrigerator at the Research Centre of Pacific Marine Resources – University of Papua (UNIPA) in Manokwari. The eggs were then shipped to ProLing Laboratory – Faculty of Fisheries and Marine Science – Bogor Agricultural University, an accredited Environmental Laboratory in Bogor, West Java for heavy metals analysis. The analysis was undertaken using the American Public Health Association (APHA) 2012 Protocols on Metals by cold-vapor atomic absorption spectrometry: Mercury (3112-B, 3030-E)[6,7]; Cadmium, Lead, Zinc, Manganese and Copper (3111-B, 3030-E) [8,7]; Arsenin (3114-B, 3030-E) [9,7]; and Iron (3500-Fe-B) [10].

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships which have, or could be perceived to have, influenced the work reported in this article.

Acknowledgments

The PI and survey team members are very grateful for the support provided by our donors, including the support of the Kaimana Government with research permit No. 005/243-SEKDA. The research (including experimental protocol and its animal use) was fully approved by the University of Papua (UNIPA) granted by Indonesia Science Authority and Ministry of Research Technology and Higher Education (Kemendiknas-Dikti) – Republic of Indonesia. We are grateful

to the lead donor: Department of Fisheries and Oceans (DFO) - Canada for providing the survey grant. Wonderful friends and committed staffs who we wish to fully acknowledge include Seiya Saleda, Piet Batubara, and boat crews at CI Kaimana. All contaminants analysis was conducted at ProLing Laboratorium – Faculty of Fisheries and Marine Science – Bogor Agricultural University in Bogor, while all field logistics details were efficiently taken care of by Heni Hokhumala.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.dib.2020.105778](https://doi.org/10.1016/j.dib.2020.105778).

References

- [1] R.F. Tapilatu, H. Wona, and P.P. Batubara., 2017. Status of sea turtle populations and its conservation at Bird's Head Seascape, Western Papua, Indonesia." *Biodiversitas Journal of Biological Diversity* 18, no. 1: 129-136. DOI: 10.13057/biodiv/d180119
- [2] B.J. Godley, D.R. Thompson, R.W. Furness., "Do heavy metal concentrations pose a threat to marine turtles from the Mediterranean Sea? *Marine Pollut. Bull.* 38 (6) (1999) 497–502.
- [3] G.F. Vazquez, M.C. Reyes, G. Fernandez., Contamination in marine turtle (*Dermochelys coriacea*) egg shells of Playon de Mexiquillo, Michoacan, Mexico, *Oceanogr. Lit. Rev.* 8 (44) (1997) 893.
- [4] BSN, Badan Standarisasi Nasional., *Batas Maksimum Cemaran Logam Berat Dalam Pangan, 7387, Standar Nasional, Indonesia, 2009*, p. 2009. SNI.
- [5] H. Sakai, H. Ichihashi, H. Sukanuma, R. Tatsukawa., "Heavy metal monitoring in sea turtles using eggs, *Marine Pollut. Bull.* 30 (5) (1995) 347–353.
- [6] Protocol 3112-B: Metals by cold-vapor atomic absorption spectrometry (<https://law.resource.org/pub/us/cfr/ibr/002/apha.method.3112.1992.pdf>) (accessed 20 March 2020).
- [7] Standard methods for the examination of water and wastewater. 3030 Preliminary Treatment of Samples (2017) (<https://www.standardmethods.org/doi/abs/10.2105/SMWW.2882.041>) (accessed 20 March 2020).
- [8] Standard Methods for the Examination of Water and Wastewater. Metals by Flame Atomic Absorption Spectrometry. <https://law.resource.org/pub/us/cfr/ibr/002/apha.method.3111.1992.pdf> (accessed 12 May 2020).
- [9] Standard methods for the examination of water and wastewater. Manual Hydride Generation Atomic Absorption Spectrometric Method (<https://law.resource.org/pub/us/cfr/ibr/002/apha.method.3114.1992.pdf>) (accessed 12 May 2020).
- [10] Standard Methods for the Examination of Water and Wastewater 22nd edition. https://beta-static.fishersci.com/content/dam/fishersci/en_US/documents/programs/scientific/technical-documents/white-papers/apha-water-testing-standard-methods-white-paper.pdf (accessed 12 May 2020).

Data on environmental contaminants in sea turtle eggs at Venu Island, Kaimana – West Papua, Indonesia

ORIGINALITY REPORT

2%

SIMILARITY INDEX

1%

INTERNET SOURCES

0%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to October University for Modern Sciences and Arts (MSA)

Student Paper

1%

2

www.mdpi.com

Internet Source

1%

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%